



US007776521B1

(12) **United States Patent**
Rota et al.

(10) **Patent No.:** **US 7,776,521 B1**
(45) **Date of Patent:** **Aug. 17, 2010**

(54) **CORONAVIRUS ISOLATED FROM HUMANS**

(75) Inventors: **Paul A. Rota**, Decatur, GA (US); **Larry J. Anderson**, Atlanta, GA (US); **William J. Bellini**, Lilburn, GA (US); **Michael D. Bowen**, Decatur, GA (US); **Cara Carthel Burns**, Avondale Estates, GA (US); **Raymond Campagnoli**, Decatur, GA (US); **Qi Chen**, Marietta, GA (US); **James A. Comer**, Decatur, GA (US); **Byron T. Cook**, Augusta, GA (US); **Shannon L. Emery**, Lusaka (ZM); **Dean D. Erdman**, Decatur, GA (US); **Cynthia S. Goldsmith**, Lilburn, GA (US); **Jeanette Guarner**, Decatur, GA (US); **Charles D. Humphrey**, Lilburn, GA (US); **Joseph P. Icenogle**, Atlanta, GA (US); **Thomas G. Ksiazek**, Lilburn, GA (US); **Richard F. Meyer**, Roswell, GA (US); **Stephan S. Monroe**, Decatur, GA (US); **William Allan Nix**, Bethlehem, GA (US); **M. Steven Oberste**, Lilburn, GA (US); **Christopher D. Paddock**, Atlanta, GA (US); **Teresa C. T. Peret**, Atlanta, GA (US); **Pierre E. Rollin**, Lilburn, GA (US); **Mark A. Pallansch**, Lilburn, GA (US); **Anthony Sanchez**, Lilburn, GA (US); **Wun-Ju Shieh**, Norcross, GA (US); **Suxiang Tong**, Alpharetta, GA (US); **Sherif R. Zaki**, Atlanta, GA (US)

(73) Assignee: **The United States of America as represented by the Secretary of the Department of Health and Human Services, Centers for Disease Control and Prevention**, Washington, DC (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 96 days.

(21) Appl. No.: **11/748,359**

(22) Filed: **May 14, 2007**

Related U.S. Application Data

(62) Division of application No. 10/822,904, filed on Apr. 12, 2004, now Pat. No. 7,220,852.

(60) Provisional application No. 60/465,927, filed on Apr. 25, 2003.

(51) **Int. Cl.**
C12Q 1/70 (2006.01)
C07H 21/00 (2006.01)
C12N 15/50 (2006.01)

(52) **U.S. Cl.** **435/5; 536/24.32; 536/24.33**

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

2005/0181357 A1* 8/2005 Peiris et al. 435/5

FOREIGN PATENT DOCUMENTS

WO WO 2004/092360 10/2004

OTHER PUBLICATIONS

Ksiazek et al (New England Journal of Medicine 348:1953-1966, published online Apr. 10, 2003).*

Gut et al (Journal of Virological Methods 77:37-46, 1999).*

Neilan et al (Nucleic Acids Research 25:2938-9, 1997).*

Peiris et al (Lancet, 361:1319-1325, published online Apr. 8, 2003).*

Genbank Accession No. AY274119.1 GI:29826276 (Apr. 14, 2003).*

SARS-associated Coronavirus. Genomic Sequence Availability. [online] [retrieved on Aug. 8, 2005]. Retrieved from the Internet <URL: <http://www.bcgsc.ca/bioinfo/SARS>>.*

"Update: Outbreak of Severe Acute Respiratory Syndrome—Worldwide, 2003," *MMWR Weekly* 52:241-248, 2003.

Emery et al., "Real-Time Reverse Transcription-Polymerase Chain Reaction Assay for SARS-Associated Coronavirus," *Emerg. Infect. Diseases* 10:311-316, 2004.

Goldsmith et al., "Ultrastructural Characterization of SARS Coronavirus," *Emerg. Infect. Diseases* 10:320-326, 2004.

Ksiazek et al., "A Novel Coronavirus Associated with Severe Acute Respiratory Syndrome," *N. Engl. J. Med.* 348:1953-1966, 2003.

Luo and Luo, "Initial SARS Coronavirus Genome Sequence Analysis Using a Bioinformatics Platform," *2nd Asia-Pacific Bioinformatics Conference (APBC2004)*, Dunedin, New Zealand, 2004.

Marra et al., "The Genome Sequence of the SARS-Associated Coronavirus," *Science* 300:1393-1404, 2003.

Supplementary Online Material for Marra et al., www.sciencemag.org/cgi/content/full/1085953/DC1, (2003).

Rota et al., "Characterization of a Novel Coronavirus Associated with Severe Acute Respiratory Syndrome," *Science* 300:1394-1399, 2003.

Supplementary Online Material for Rota et al., www.sciencemag.org/cgi/content/full/1085953/DC1, (2003).

GenBank Accession No. AY274119, Apr. 14, 2003.

GenBank Accession No. AY278741, Apr. 21, 2003.

GenBank Accession No. AY278554.1, Apr. 18, 2003.

GenBank Accession No. AY278491.1, Apr. 18, 2003.

GenBank Accession No. AY278487, Apr. 21, 2003.

* cited by examiner

Primary Examiner—Mary E Mosher
(74) *Attorney, Agent, or Firm*—Klarquist Sparkman, LLP

(57) **ABSTRACT**

Disclosed herein is a newly isolated human coronavirus (SARS-CoV), the causative agent of severe acute respiratory syndrome (SARS). Also provided are the nucleic acid sequence of the SARS-CoV genome and the amino acid sequences of the SARS-CoV open reading frames, as well as methods of using these molecules to detect a SARS-CoV and detect infections therewith. Immune stimulatory compositions are also provided, along with methods of their use.

7 Claims, 7 Drawing Sheets




FIG. 1A




FIG. 1B




FIG. 2A




FIG. 2B




FIG. 3




FIG. 4

FIG. 5A




FIG. 5B





FIG. 5C




FIG. 6A




FIG. 6B

FIG. 7A





FIG. 7B



CORONAVIRUS ISOLATED FROM HUMANS

PRIORITY CLAIM

This is a division of co-pending U.S. patent application Ser. No. 10/822,904, filed Apr. 12, 2004, and issued as U.S. Pat. No. 7,220,852 on May 22, 2007, which in turn claims the benefit of U.S. Provisional Patent Application No. 60/465,927 filed Apr. 25, 2003. Both applications are incorporated herein by reference in their entirety.

STATEMENT OF GOVERNMENT SUPPORT

This invention was made by the Centers for Disease Control and Prevention, an agency of the United States Government. Therefore, the U.S. Government has certain rights in this invention.

FIELD OF THE DISCLOSURE

This invention relates to a newly isolated human coronavirus. More particularly, it relates to an isolated coronavirus genome, isolated coronavirus proteins, and isolated nucleic acid molecules encoding the same. The disclosure further relates to methods of detecting a severe acute respiratory syndrome-associated coronavirus and compositions comprising immunogenic coronavirus compounds.

BACKGROUND

The coronaviruses (order Nidovirales, family Coronaviridae, genus *Coronavirus*) are a diverse group of large, enveloped, positive-stranded RNA viruses that cause respiratory and enteric diseases in humans and other animals. At approximately 30,000 nucleotides (nt), their genome is the largest found in any of the RNA viruses. Coronaviruses are spherical, 100-160 nm in diameter with 20-40 nm complex club shaped surface projections surrounding the periphery. Coronaviruses share common structural proteins including a spike protein (S), membrane protein (M), envelope protein (E), and, in a subset of coronaviruses, a hemagglutinin-esterase protein (HE). The S protein, a glycoprotein which protrudes from the virus membrane, is involved in host cell receptor binding and is a target for neutralizing antibodies. The E and M proteins are involved in virion formation and release from the host cell. Coronavirus particles are found within the cisternae of the rough endoplasmic reticulum and in vesicles of infected host cells where virions are assembled. The coronavirus genome consists of two open reading frames (ORF1a and ORF1b) yielding an RNA polymerase and a nested set of subgenomic mRNAs encoding structural and nonstructural proteins, including the S, E, M, and nucleocapsid (N) proteins. The genus *Coronavirus* includes at least 13 species which have been subdivided into at least three groups (groups I, II, and III) on the basis of serological and genetic properties (deVries et al., *Sem. Virol.* 8:33-47, 1997; Fields et al. eds. *Fields Virology*, 3rd edition, Raven Press, Philadelphia, 1323-1341, 1996; Mahey and Collier eds. *Microbiology and Microbial Infections*, Volume 1 Virology, 9th edition, Oxford University Press, 463-479, 1998).

The three known groups of coronavirus are associated with a variety of diseases of humans and domestic animals (for example, cattle, pigs, cats, dogs, rodents, and birds), including gastroenteritis and upper and lower respiratory tract disease. Known coronaviruses include human Coronavirus 229E (HCoV-229E), canine coronavirus (CCoV), feline infectious peritonitis virus (FIPV), porcine transmissible gas-

troenteritis virus (TGEV), porcine epidemic diarrhea virus (PEDV), human coronavirus OC43 (HCoV-OC43), bovine coronavirus (BCoV), porcine hemagglutinating encephalomyelitis virus (HEV), rat sialodacryoadenitis virus (SDAV), mouse hepatitis virus (MHV), turkey coronavirus (TCoV), and avian infectious bronchitis virus (IBV-Avian) (Fields et al. eds. *Fields Virology*, 3rd edition, Raven Press, Philadelphia, 1323-1341, 1996; Mahey and Collier eds. *Microbiology and Microbial Infections*, Volume 1 Virology, 9th edition, Oxford University Press, 463-479, 1998).

Coronavirus infections are generally host specific with respect to infectivity and clinical symptoms. Coronaviruses further exhibit marked tissue tropism; infection in the incorrect host species or tissue type may result in an abortive infection, mutant virus production and altered virulence. Coronaviruses generally do not grow well in cell culture, and animal models for human *coronavirus* infection are lacking. Therefore, little is known about them (Fields et al. eds. *Fields Virology*, 3rd edition, Raven Press, Philadelphia, 1323-1341, 1996). The known human coronaviruses are notably fastidious in cell culture, preferring select cell lines, organ culture, or suckling mice for propagation. Coronaviruses grown in cell culture exhibit varying degrees of virulence and/or cytopathic effect (CPE) depending on the host cell type and culture conditions. The only human or animal coronavirus which has been shown to grow in Vero E6 cells is PEDV, and it requires the addition of trypsin to culture medium for growth in Vero E6 cells. Moreover, PEDV adapted to Vero E6 cell culture results in a strikingly different CPE, with cytoplasmic vacuoles and the formation of large syncytia (Hofmann and Wyler, *J. Clin. Micro.* 26:2235-39, 1988; Kusanagi et al., *J. Vet. Med. Sci.* 554:313-18, 1991).

Coronavirus have not previously been known to cause severe disease in humans, but have been identified as a major cause of upper respiratory tract illness, including the common cold. Repeat infections in humans are common within and across serotype, suggesting that immune response to coronavirus infection in humans is either incomplete or short lived. Coronavirus infection in animals can cause severe enteric or respiratory disease. Vaccination has been used successfully to prevent and control some coronavirus infections in animals. The ability of animal-specific coronaviruses to cause severe disease raises the possibility that coronavirus could also cause more severe disease in humans (Fields et al. eds. *Fields Virology*, 3rd edition, Raven Press, Philadelphia, 1323-1341, 1996; Mahey and Collier eds. *Microbiology and Microbial Infections*, Volume 1 Virology, 9th edition, Oxford University Press, 463-479, 1998).

In late 2002, cases of life-threatening respiratory disease with no identifiable etiology were reported from Guangdong Province, China, followed by reports from Vietnam, Canada, and Hong Kong of severe febrile respiratory illness that spread to household members and health care workers. The syndrome was designated "severe acute respiratory syndrome" (SARS) in February 2003 by the Centers for Disease Control and Prevention (*MMWR*, 52:241-48, 2003).

Past efforts to develop rapid diagnostics and vaccines for coronavirus infection in humans have been hampered by a lack of appropriate research models and the moderate course of disease in humans. Therefore, a need for rapid diagnostic tests and vaccines exists.

SUMMARY OF THE DISCLOSURE

A newly isolated human coronavirus has been identified as the causative agent of SARS, and is termed SARS-CoV. The

nucleic acid sequence of the SARS-CoV genome and the amino acid sequences of the SARS-CoV open reading frames are provided herein.

This disclosure provides methods and compositions useful in detecting the presence of a SARS-CoV nucleic acid in a sample and/or diagnosing a SARS-CoV infection in a subject. Also provided are methods and compositions useful in detecting the presence of a SARS-CoV antigen or antibody in a sample and/or diagnosing a SARS-CoV infection in a subject.

The foregoing and other features and advantages will become more apparent from the following detailed description of several embodiments, which proceeds with reference to the accompanying figures.

BRIEF DESCRIPTION OF THE FIGURES

FIGS. 1A-B are photomicrographs illustrating typical early cytopathic effects seen with coronavirus isolates and serum from SARS patients. FIG. 1A is a photomicrograph of Vero E6 cells inoculated with an oropharyngeal specimen from a SARS patient (x40). FIG. 1B is a photomicrograph of infected Vero E6 cells reacting with the serum of a convalescent SARS patient in an indirect fluorescent antibody (IFA) assay (x400).

FIGS. 2A-B are electronmicrographs illustrating ultrastructural characteristics of the SARS-associated coronavirus (SARS-CoV). FIG. 2A is a thin-section electron-microscopical view of viral nucleocapsids aligned along the membrane of the rough endoplasmic reticulum (arrow) as particles bud into the cisternae. Enveloped virions have surface projections (arrowhead) and an electron-lucent center. Directly under the viral envelope lies a characteristic ring formed by the helical nucleocapsid, often seen in cross-section. FIG. 2B is a negative stain (methylamine tungstate) electronmicrograph showing stain-penetrated coronavirus particle with the typical internal helical nucleocapsid-like structure and club-shaped surface projections surrounding the periphery of the particle. Bars: 100 nm.

FIG. 3 is an estimated maximum parsimony tree illustrating putative phylogenetic relationships between SARS-CoV and other human and animal coronaviruses. Phylogenetic relationships are based on sequence alignment of 405 nucleotides of the coronavirus polymerase gene ORF1b (nucleic acid 15,173 to 15,578 of SEQ ID NO: 1). The three major coronavirus antigenic groups (I, II and III), represented by HCoV-229E, CCoV, FIPV, TGEV, PEDV, HCoV-OC43, BCoV, HEV, SDAV, MHV, TCoV, and IBV-Avian, are shown shaded. Bootstrap values (100 replicates) obtained from a 50% majority rule consensus tree are plotted at the main internal branches of the phylogram. Branch lengths are proportionate to nucleotide differences.

FIG. 4 is a pictorial representation of neighbor joining trees illustrating putative phylogenetic relationships between SARS-CoV and other human and animal coronaviruses. Amino acid sequences of the indicated SARS-CoV proteins were compared with those from reference viruses representing each species in the three groups of coronaviruses for which complete genomic sequence information was available [group 1: HCoV-229E (AF304460); PEDV (AF353511); TGEV (AJ271965); group 2: BCoV (AF220295); MHV (AF201929); group 3: infectious bronchitis virus (M95169)]. Sequences for representative strains of other coronavirus species, for which partial sequence information was available, were included for some of the structural protein comparisons [group 1: CCoV (D13096); FCoV (AY204704); porcine respiratory coronavirus (Z24675); group 2: HCoV-OC43 (M76373, L14643, M93390); HEV (AY078417); rat coro-

navirus (AF207551)]. Sequence alignments and neighbor joining trees were generated by using Clustalx 1.83 with the Gonnet protein comparison matrix. The resulting trees were adjusted for final output using treeotool 2.0.1.

FIGS. 5A-C are photomicrographs illustrating diffuse alveolar damage in a patient with SARS (FIGS. 5A-B), and immunohistochemical staining of SARS-CoV-infected Vero E6 cells (FIG. 5C). FIG. 5A is a photomicrograph of lung tissue from a SARS patient (x50). Diffuse alveolar damage, abundant foamy macrophages and multinucleated syncytial cells are present; hematoxylin and eosin stain was used. FIG. 5B is a higher magnification photomicrograph of lung tissue from the same SARS patient (x250). Syncytial cells show no conspicuous viral inclusions. FIG. 5C is a photomicrograph of immunohistochemically stained SARS-CoV-infected cells (x250). Membranous and cytoplasmic immunostaining of individual and syncytial Vero E6 cells was achieved using feline anti-FIPV-1 ascitic fluid. Immunoalkaline phosphatase with naphthol-fast red substrate and hematoxylin counter stain was used.

FIGS. 6A-B are electronmicrographs illustrating ultrastructural characteristics of a coronavirus-infected cell in bronchoalveolar lavage (BAL) from a SARS patient. FIG. 6A is an electron micrograph of a coronavirus-infected cell. Numerous intracellular and extracellular particles are present; virions are indicated by the arrowheads. FIG. 6B is a higher magnification electronmicrograph of the area seen at the arrow in FIG. 6A (rotated clockwise approximately 90°). Bars: FIG. 6A, 1 µm; FIG. 6B, 100 nm.

FIGS. 7A-C illustrate the organization of the SARS-CoV genome. FIG. 7A is a diagram of the overall organization of the 29,727-nt SARS-CoV genomic RNA. The 72-nt leader sequence is represented as a small rectangle at the left-most end. ORFs1a and 1b, encoding the nonstructural polyproteins, and those ORFs encoding the S, E, M, and N structural proteins, are indicated. Vertical position of the boxes indicates the phase of the reading frame (phase 1 for proteins above the line, phase two for proteins on the line and phase three for proteins below the line). FIG. 7B is an expanded view of the structural protein encoding region and predicted mRNA transcripts. Known structural protein encoding regions (dark grey boxes) and regions and reading frames for potential products X1-X5 (light gray boxes) are indicated. Lengths and map locations of the 3'-coterminal mRNAs expressed by the SARS-CoV are indicated, as predicted by identification of conserved transcriptional regulatory sequences. FIG. 7C is a digitized image of a nylon membrane showing Northern blot analysis of SARS-CoV mRNAs. Poly (A)⁺ RNA from infected Vero E6 cells was separated on a formaldehyde-agarose gel, transferred to a nylon membrane, and hybridized with a digoxigenin-labeled riboprobe overlapping the 3'-untranslated region. Signals were visualized by chemiluminescence. Sizes of the SARS-CoV mRNAs were calculated by extrapolation from a log-linear fit of the molecular mass marker. Lane 1, SARS-CoV mRNA; lane 2, Vero E6 cell mRNA; lane 3, molecular mass marker, sizes in kB.

SEQUENCE LISTING

The nucleic and amino acid sequences listed in the accompanying sequence listing are shown using standard letter abbreviations for nucleotide bases, and three letter code for amino acids, as defined in 37 C.F.R. 1.822. Only one strand of each nucleic acid sequence is shown, but the complementary strand is understood as included by any reference to the displayed strand. In the accompanying sequence listing:

5

SEQ ID NO: 1 shows the nucleic acid sequence of the SARS-CoV genome.

SEQ ID NO: 2 shows the amino acid sequence of the SARS-CoV polyprotein 1a (encoded by nucleic acid 265 to nucleic acid 13,398 of SEQ ID NO: 1).

SEQ ID NO: 3 shows the amino acid sequence of the SARS-CoV polyprotein 1b (encoded by nucleic acid 13,398 to 21,482 of SEQ ID NO: 1).

SEQ ID NO: 4 shows the amino acid sequence of the SARS-CoV S protein (encoded by nucleic acid 21,492 to 25,256 of SEQ ID NO: 1).

SEQ ID NO: 5 shows the amino acid sequence of the SARS-CoV X1 protein (encoded by nucleic acid 25,268 to 26,089 of SEQ ID NO: 1).

SEQ ID NO: 6 shows the amino acid sequence of the SARS-CoV X2 protein (encoded by nucleic acid 25,689 to 26,150 of SEQ ID NO: 1).

SEQ ID NO: 7 shows the amino acid sequence of the SARS-CoV E protein (encoded by nucleic acid 26,117 to 26,344 of SEQ ID NO: 1).

SEQ ID NO: 8 shows the amino acid sequence of the SARS-CoV M protein (encoded by nucleic acid 26,398 to 27,060 of SEQ ID NO: 1).

SEQ ID NO: 9 shows the amino acid sequence of the SARS-CoV X3 protein (encoded by nucleic acid 27,074 to 27,262 of SEQ ID NO: 1).

SEQ ID NO: 10 shows the amino acid sequence of the SARS-CoV X4 protein (encoded by nucleic acid 27,273 to 27,638 of SEQ ID NO: 1).

SEQ ID NO: 11 shows the amino acid sequence of the SARS-CoV X5 protein (encoded by nucleic acid 27,864 to 28,115 of SEQ ID NO: 1).

SEQ ID NO: 12 shows the amino acid sequence of the SARS-CoV N protein (encoded by nucleic acid 28,120 to 29,385 of SEQ ID NO: 1).

SEQ ID NOS: 13-15 show the nucleic acid sequence of several SARS-CoV-specific oligonucleotide primers.

SEQ ID NOS: 16-33 show the nucleic acid sequence of several oligonucleotide primers/probes used for real-time reverse transcription-polymerase chain reaction (RT-PCR) SARS-CoV assays.

SEQ ID NOS: 34-35 show the nucleic acid sequence of two degenerate primers designed to anneal to sites encoding conserved coronavirus amino acid motifs.

SEQ ID NOS: 36-38 show the nucleic acid sequence of several oligonucleotide primers/probes used as controls in real-time RT-PCR assays.

DETAILED DESCRIPTION OF SEVERAL EMBODIMENTS

I. Abbreviations

BAL: bronchoalveolar lavage
 CPE: cytopathic effect
 E: coronavirus transmembrane protein
 ELISA: enzyme-linked immunosorbent assay
 HE: coronavirus hemagglutinin-esterase protein
 IFA: indirect fluorescent antibody
 M: coronavirus membrane protein
 N: coronavirus nucleoprotein
 ORF: open reading frame
 PCR: polymerase chain reaction
 RACE: 5' rapid amplification of cDNA ends
 RT-PCR: reverse transcription-polymerase chain reaction
 S: coronavirus spike protein
 SARS: severe acute respiratory syndrome

6

SARS-CoV: severe acute respiratory syndrome-associated coronavirus

TRS: transcriptional regulatory sequence

II. Terms

Unless otherwise noted, technical terms are used according to conventional usage. Definitions of common terms in molecular biology may be found in Benjamin Lewin, *Genes VII*, published by Oxford University Press, 2000 (ISBN 019879276X); Kendrew et al. (eds.), *The Encyclopedia of Molecular Biology*, published by Blackwell Publishers, 1994 (ISBN 0632021829); and Robert A. Meyers (ed.), *Molecular Biology and Biotechnology: a Comprehensive Desk Reference*, published by Wiley, John & Sons, Inc., 1995 (ISBN 0471186341); and other similar references.

As used herein, the singular terms "a," "an," and "the" include plural referents unless context clearly indicates otherwise. Similarly, the word "or" is intended to include "and" unless the context clearly indicates otherwise. Also, as used herein, the term "comprises" means "includes." Hence "comprising A or B" means including A, B, or A and B. It is further to be understood that all nucleotide sizes or amino acid sizes, and all molecular weight or molecular mass values, given for nucleic acids or polypeptides are approximate, and are provided for description. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including explanations of terms, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

In order to facilitate review of the various embodiments of this disclosure, the following explanations of specific terms are provided:

Adjuvant: A substance that non-specifically enhances the immune response to an antigen. Development of vaccine adjuvants for use in humans is reviewed in Singh et al. (*Nat. Biotechnol.* 17:1075-1081, 1999), which discloses that, at the time of its publication, aluminum salts and the MF59 micro-emulsion are the only vaccine adjuvants approved for human use.

Amplification: Amplification of a nucleic acid molecule (e.g., a DNA or RNA molecule) refers to use of a laboratory technique that increases the number of copies of a nucleic acid molecule in a sample. An example of amplification is the polymerase chain reaction (PCR), in which a sample is contacted with a pair of oligonucleotide primers under conditions that allow for the hybridization of the primers to a nucleic acid template in the sample. The primers are extended under suitable conditions, dissociated from the template, re-annealed, extended, and dissociated to amplify the number of copies of the nucleic acid. The product of amplification can be characterized by such techniques as electrophoresis, restriction endonuclease cleavage patterns, oligonucleotide hybridization or ligation, and/or nucleic acid sequencing.

Other examples of amplification methods include strand displacement amplification, as disclosed in U.S. Pat. No. 5,744,311; transcription-free isothermal amplification, as disclosed in U.S. Pat. No. 6,033,881; repair chain reaction amplification, as disclosed in WO 90/01069; ligase chain reaction amplification, as disclosed in EP-A-320,308; gap filling ligase chain reaction amplification, as disclosed in U.S. Pat. No. 5,427,930; and NASBA™ RNA transcription-free amplification, as disclosed in U.S. Pat. No. 6,025,134. An

amplification method can be modified, including for example by additional steps or coupling the amplification with another protocol.

Animal: Living multi-cellular vertebrate organisms, a category that includes, for example, mammals and birds. The term mammal includes both human and non-human mammals. Similarly, the term "subject" includes both human and veterinary subjects, for example, humans, non-human primates, dogs, cats, horses, and cows.

Antibody: A protein (or protein complex) that includes one or more polypeptides substantially encoded by immunoglobulin genes or fragments of immunoglobulin genes. The recognized immunoglobulin genes include the kappa, lambda, alpha, gamma, delta, epsilon, and mu constant region genes, as well as the myriad immunoglobulin variable region genes. Light chains are classified as either kappa or lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, which in turn define the immunoglobulin classes, IgG, IgM, IgA, IgD and IgE, respectively.

The basic immunoglobulin (antibody) structural unit is generally a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one "light" (about 25 kDa) and one "heavy" (about 50-70 kDa) chain. The N-terminus of each chain defines a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The terms "variable light chain" (V_L) and "variable heavy chain" (V_H) refer, respectively, to these light and heavy chains

As used herein, the term "antibodies" includes intact immunoglobulins as well as a number of well-characterized fragments. For instance, Fabs, Fvs, and single-chain Fvs (SCFvs) that bind to target protein (or epitope within a protein or fusion protein) would also be specific binding agents for that protein (or epitope). These antibody fragments are defined as follows: (1) Fab, the fragment which contains a monovalent antigen-binding fragment of an antibody molecule produced by digestion of whole antibody with the enzyme papain to yield an intact light chain and a portion of one heavy chain; (2) Fab', the fragment of an antibody molecule obtained by treating whole antibody with pepsin, followed by reduction, to yield an intact light chain and a portion of the heavy chain; two Fab' fragments are obtained per antibody molecule; (3) $(Fab')_2$, the fragment of the antibody obtained by treating whole antibody with the enzyme pepsin without subsequent reduction; (4) $F(ab')_2$, a dimer of two Fab' fragments held together by two disulfide bonds; (5) Fv, a genetically engineered fragment containing the variable region of the light chain and the variable region of the heavy chain expressed as two chains; and (6) single chain antibody, a genetically engineered molecule containing the variable region of the light chain, the variable region of the heavy chain, linked by a suitable polypeptide linker as a genetically fused single chain molecule. Methods of making these fragments are routine (see, for example, Harlow and Lane, *Using Antibodies: A Laboratory Manual*, CSHL, New York, 1999).

Antibodies for use in the methods and devices of this disclosure can be monoclonal or polyclonal. Merely by way of example, monoclonal antibodies can be prepared from murine hybridomas according to the classical method of Kohler and Milstein (*Nature* 256:495-97, 1975) or derivative methods thereof. Detailed procedures for monoclonal antibody production are described in Harlow and Lane, *Using Antibodies: A Laboratory Manual*, CSHL, New York, 1999.

Antigen: A compound, composition, or substance that can stimulate the production of antibodies or a T-cell response in an animal, including compositions that are injected or absorbed into an animal. An antigen reacts with the products

of specific humoral or cellular immunity, including those induced by heterologous immunogens. In one embodiment, an antigen is a coronavirus antigen.

Binding or Stable Binding: An oligonucleotide binds or stably binds to a target nucleic acid if a sufficient amount of the oligonucleotide forms base pairs or is hybridized to its target nucleic acid, to permit detection of that binding. Binding can be detected by either physical or functional properties of the target:oligonucleotide complex. Binding between a target and an oligonucleotide can be detected by any procedure known to one skilled in the art, including functional or physical binding assays. Binding can be detected functionally by determining whether binding has an observable effect upon a biosynthetic process such as expression of a gene, DNA replication, transcription, translation, and the like.

Physical methods of detecting the binding of complementary strands of DNA or RNA are well known in the art, and include such methods as DNase I or chemical footprinting, gel shift and affinity cleavage assays, Northern blotting, Southern blotting, dot blotting, and light absorption detection procedures. For example, a method which is widely used, because it is so simple and reliable, involves observing a change in light absorption of a solution containing an oligonucleotide (or an analog) and a target nucleic acid at 220 to 300 nm as the temperature is slowly increased. If the oligonucleotide or analog has bound to its target, there is a sudden increase in absorption at a characteristic temperature as the oligonucleotide (or analog) and target dissociate or melt.

The binding between an oligomer and its target nucleic acid is frequently characterized by the temperature (T_m) at which 50% of the oligomer is melted from its target. A higher T_m means a stronger or more stable complex relative to a complex with a lower T_m .

cDNA (complementary DNA): A piece of DNA lacking internal, non-coding segments (introns) and regulatory sequences that determine transcription. cDNA is synthesized in the laboratory by reverse transcription from messenger RNA extracted from cells.

Electrophoresis: Electrophoresis refers to the migration of charged solutes or particles in a liquid medium under the influence of an electric field. Electrophoretic separations are widely used for analysis of macromolecules. Of particular importance is the identification of proteins and nucleic acid sequences. Such separations can be based on differences in size and/or charge. Nucleotide sequences have a uniform charge and are therefore separated based on differences in size. Electrophoresis can be performed in an unsupported liquid medium (for example, capillary electrophoresis), but more commonly the liquid medium travels through a solid supporting medium. The most widely used supporting media are gels, for example, polyacrylamide and agarose gels.

Sieving gels (for example, agarose) impede the flow of molecules. The pore size of the gel determines the size of a molecule that can flow freely through the gel. The amount of time to travel through the gel increases as the size of the molecule increases. As a result, small molecules travel through the gel more quickly than large molecules and thus progress further from the sample application area than larger molecules, in a given time period. Such gels are used for size-based separations of nucleotide sequences.

Fragments of linear DNA migrate through agarose gels with a mobility that is inversely proportional to the \log_{10} of their molecular weight. By using gels with different concentrations of agarose, different sizes of DNA fragments can be resolved. Higher concentrations of agarose facilitate separation of small DNAs, while low agarose concentrations allow resolution of larger DNAs.

Hybridization: Oligonucleotides and their analogs hybridize by hydrogen bonding, which includes Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding, between complementary bases. Generally, nucleic acid consists of nitrogenous bases that are either pyrimidines (cytosine (C), uracil (U), and thymine (T)) or purines (adenine (A) and guanine (G)). These nitrogenous bases form hydrogen bonds between a pyrimidine and a purine, and the bonding of the pyrimidine to the purine is referred to as "base pairing." More specifically, A will hydrogen bond to T or U, and G will bond to C. "Complementary" refers to the base pairing that occurs between two distinct nucleic acid sequences or two distinct regions of the same nucleic acid sequence.

"Specifically hybridizable" and "specifically complementary" are terms that indicate a sufficient degree of complementarity such that stable and specific binding occurs between the oligonucleotide (or its analog) and the DNA or RNA target. The oligonucleotide or oligonucleotide analog need not be 100% complementary to its target sequence to be specifically hybridizable. An oligonucleotide or analog is specifically hybridizable when binding of the oligonucleotide or analog to the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA, and there is a sufficient degree of complementarity to avoid non-specific binding of the oligonucleotide or analog to non-target sequences under conditions where specific binding is desired, for example under physiological conditions in the case of in vivo assays or systems. Such binding is referred to as specific hybridization.

Hybridization conditions resulting in particular degrees of stringency will vary depending upon the nature of the hybridization method of choice and the composition and length of the hybridizing nucleic acid sequences. Generally, the temperature of hybridization and the ionic strength (especially the Na⁺ and/or Mg⁺⁺ concentration) of the hybridization buffer will determine the stringency of hybridization, though wash times also influence stringency. Calculations regarding hybridization conditions required for attaining particular degrees of stringency are discussed by Sambrook et al. (ed.), *Molecular Cloning: A Laboratory Manual*, 2nd ed., vol. 1-3, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989, chapters 9 and 11; and Ausubel et al. *Short Protocols in Molecular Biology*, 4th ed., John Wiley & Sons, Inc., 1999.

For purposes of the present disclosure, "stringent conditions" encompass conditions under which hybridization will only occur if there is less than 25% mismatch between the hybridization molecule and the target sequence. "Stringent conditions" may be broken down into particular levels of stringency for more precise definition. Thus, as used herein, "moderate stringency" conditions are those under which molecules with more than 25% sequence mismatch will not hybridize; conditions of "medium stringency" are those under which molecules with more than 15% mismatch will not hybridize, and conditions of "high stringency" are those under which sequences with more than 10% mismatch will not hybridize. Conditions of "very high stringency" are those under which sequences with more than 6% mismatch will not hybridize.

Immune Stimulatory Composition: A term used herein to mean a composition useful for stimulating or eliciting a specific immune response (or immunogenic response) in a vertebrate. In some embodiments, the immunogenic response is protective or provides protective immunity, in that it enables

the vertebrate animal to better resist infection with or disease progression from the organism against which the vaccine is directed.

Without wishing to be bound by a specific theory, it is believed that an immunogenic response may arise from the generation of an antibody specific to one or more of the epitopes provided in the immune stimulatory composition. Alternatively, the response may comprise a T-helper or cytotoxic cell-based response to one or more of the epitopes provided in the immune stimulatory composition. All three of these responses may originate from naïve or memory cells. One specific example of a type of immune stimulatory composition is a vaccine.

In some embodiments, an "effective amount" or "immune-stimulatory amount" of an immune stimulatory composition is an amount which, when administered to a subject, is sufficient to engender a detectable immune response. Such a response may comprise, for instance, generation of an antibody specific to one or more of the epitopes provided in the immune stimulatory composition. Alternatively, the response may comprise a T-helper or CTL-based response to one or more of the epitopes provided in the immune stimulatory composition. All three of these responses may originate from naïve or memory cells. In other embodiments, a "protective effective amount" of an immune stimulatory composition is an amount which, when administered to a subject, is sufficient to confer protective immunity upon the subject.

Inhibiting or Treating a Disease: Inhibiting the full development of a disease or condition, for example, in a subject who is at risk for a disease such as SARS. "Treatment" refers to a therapeutic intervention that ameliorates a sign or symptom of a disease or pathological condition after it has begun to develop. As used herein, the term "ameliorating," with reference to a disease, pathological condition or symptom, refers to any observable beneficial effect of the treatment. The beneficial effect can be evidenced, for example, by a delayed onset of clinical symptoms of the disease in a susceptible subject, a reduction in severity of some or all clinical symptoms of the disease, a slower progression of the disease, a reduction in the number of relapses of the disease, an improvement in the overall health or well-being of the subject, or by other parameters well known in the art that are specific to the particular disease.

Isolated: An "isolated" microorganism (such as a virus, bacterium, fungus, or protozoan) has been substantially separated or purified away from microorganisms of different types, strains, or species. Microorganisms can be isolated by a variety of techniques, including serial dilution and culturing.

An "isolated" biological component (such as a nucleic acid molecule, protein or organelle) has been substantially separated or purified away from other biological components in the cell of the organism in which the component naturally occurs, such as other chromosomal and extra-chromosomal DNA and RNA, proteins, and organelles. Nucleic acids and proteins that have been "isolated" include nucleic acids and proteins purified by standard purification methods. The term also embraces nucleic acids and proteins prepared by recombinant expression in a host cell, as well as chemically synthesized nucleic acids or proteins, or fragments thereof.

Label: A detectable compound or composition that is conjugated directly or indirectly to another molecule to facilitate detection of that molecule. Specific, non-limiting examples of labels include fluorescent tags, enzymatic linkages, and radioactive isotopes.

Nucleic Acid Molecule: A polymeric form of nucleotides, which may include both sense and anti-sense strands of RNA,

cDNA, genomic DNA, and synthetic forms and mixed polymers of the above. A nucleotide refers to a ribonucleotide, deoxynucleotide or a modified form of either type of nucleotide. A "nucleic acid molecule" as used herein is synonymous with "nucleic acid" and "polynucleotide." A nucleic acid molecule is usually at least 10 bases in length, unless otherwise specified. The term includes single- and double-stranded forms of DNA. A polynucleotide may include either or both naturally occurring and modified nucleotides linked together by naturally occurring and/or non-naturally occurring nucleotide linkages.

Oligonucleotide: A nucleic acid molecule generally comprising a length of 300 bases or fewer. The term often refers to single-stranded deoxyribonucleotides, but it can refer as well to single- or double-stranded ribonucleotides, RNA:DNA hybrids and double-stranded DNAs, among others. The term "oligonucleotide" also includes oligonucleosides (that is, an oligonucleotide minus the phosphate) and any other organic base polymer. In some examples, oligonucleotides are about 10 to about 90 bases in length, for example, 12, 13, 14, 15, 16, 17, 18, 19 or 20 bases in length. Other oligonucleotides are about 25, about 30, about 35, about 40, about 45, about 50, about 55, about 60 bases, about 65 bases, about 70 bases, about 75 bases or about 80 bases in length. Oligonucleotides may be single-stranded, for example, for use as probes or primers, or may be double-stranded, for example, for use in the construction of a mutant gene. Oligonucleotides can be either sense or anti-sense oligonucleotides. An oligonucleotide can be modified as discussed above in reference to nucleic acid molecules. Oligonucleotides can be obtained from existing nucleic acid sources (for example, genomic or cDNA), but can also be synthetic (for example, produced by laboratory or in vitro oligonucleotide synthesis).

Open Reading Frame (ORF): A series of nucleotide triplets (codons) coding for amino acids without any internal termination codons. These sequences are usually translatable into a peptide/polypeptide/protein/polypeptide.

Operably Linked: A first nucleic acid sequence is operably linked with a second nucleic acid sequence when the first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Generally, operably linked DNA sequences are contiguous and, where necessary to join two protein coding regions, in the same reading frame. If introns are present, the operably linked DNA sequences may not be contiguous.

Pharmaceutically Acceptable Carriers: The pharmaceutically acceptable carriers useful in this disclosure are conventional. *Remington's Pharmaceutical Sciences*, by E. W. Martin, Mack Publishing Co., Easton, Pa., 15th Edition (1975), describes compositions and formulations suitable for pharmaceutical delivery of one or more therapeutic compounds or molecules, such as one or more SARS-CoV nucleic acid molecules, proteins or antibodies that bind these proteins, and additional pharmaceutical agents.

In general, the nature of the carrier will depend on the particular mode of administration being employed. For instance, parenteral formulations usually comprise injectable fluids that include pharmaceutically and physiologically acceptable fluids such as water, physiological saline, balanced salt solutions, aqueous dextrose, glycerol or the like as a vehicle. For solid compositions (for example, powder, pill, tablet, or capsule forms), conventional non-toxic solid carriers can include, for example, pharmaceutical grades of mannitol, lactose, starch, or magnesium stearate. In addition to biologically-neutral carriers, pharmaceutical compositions to

be administered can contain minor amounts of non-toxic auxiliary substances, such as wetting or emulsifying agents, preservatives, and pH buffering agents and the like, for example sodium acetate or sorbitan monolaurate.

Polypeptide: A polymer in which the monomers are amino acid residues which are joined together through amide bonds. When the amino acids are alpha-amino acids, either the L-optical isomer or the D-optical isomer can be used, the L-isomers being preferred. The terms "polypeptide" or "protein" as used herein are intended to encompass any amino acid sequence and include modified sequences such as glycoproteins. The term "polypeptide" is specifically intended to cover naturally occurring proteins, as well as those which are recombinantly or synthetically produced.

Conservative amino acid substitutions are those substitutions that, when made, least interfere with the properties of the original protein, that is, the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. Examples of conservative substitutions are shown below.

Original Residue	Conservative Substitutions
Ala	Ser
Arg	Lys
Asn	Gln, His
Asp	Glu
Cys	Ser
Gln	Asn
Glu	Asp
His	Asn; Gln
Ile	Leu, Val
Leu	Ile; Val
Lys	Arg; Gln; Glu
Met	Leu; Ile
Phe	Met; Leu; Tyr
Ser	Thr
Thr	Ser
Trp	Tyr
Tyr	Trp; Phe
Val	Ile; Leu

Conservative substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain.

The substitutions which in general are expected to produce the greatest changes in protein properties will be non-conservative, for instance changes in which (a) a hydrophilic residue, for example, seryl or threonyl, is substituted for (or by) a hydrophobic residue, for example, leucyl, isoleucyl, phenylalanyl, valyl or alanyl; (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue having an electropositive side chain, for example, lysyl, arginyl, or histadyl, is substituted for (or by) an electronegative residue, for example, glutamyl or aspartyl; or (d) a residue having a bulky side chain, for example, phenylalanine, is substituted for (or by) one not having a side chain, for example, glycine.

Probes and Primers: A probe comprises an isolated nucleic acid attached to a detectable label or other reporter molecule. Typical labels include radioactive isotopes, enzyme substrates, co-factors, ligands, chemiluminescent or fluorescent agents, haptens, and enzymes. Methods for labeling and guidance in the choice of labels appropriate for various purposes are discussed, for example, in Sambrook et al. (ed.), *Molecular Cloning: A Laboratory Manual*, 2nd ed., vol. 1-3, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.,

1989 and Ausubel et al. *Short Protocols in Molecular Biology*, 4th ed., John Wiley & Sons, Inc., 1999.

Primers are short nucleic acid molecules, for instance DNA oligonucleotides 10 nucleotides or more in length, for example that hybridize to contiguous complementary nucleotides or a sequence to be amplified. Longer DNA oligonucleotides may be about 15, 20, 25, 30 or 50 nucleotides or more in length. Primers can be annealed to a complementary target DNA strand by nucleic acid hybridization to form a hybrid between the primer and the target DNA strand, and then the primer extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification of a nucleic acid sequence, for example, by the PCR or other nucleic-acid amplification methods known in the art, as described above.

Methods for preparing and using nucleic acid probes and primers are described, for example, in Sambrook et al. (ed.), *Molecular Cloning: A Laboratory Manual*, 2nd ed., vol. 1-3, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989; Ausubel et al. *Short Protocols in Molecular Biology*, 4th ed., John Wiley & Sons, Inc., 1999; and Innis et al. *PCR Protocols, A Guide to Methods and Applications*, Academic Press, Inc., San Diego, Calif., 1990. Amplification primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, © 1991, Whitehead Institute for Biomedical Research, Cambridge, Mass.). One of ordinary skill in the art will appreciate that the specificity of a particular probe or primer increases with its length. Thus, in order to obtain greater specificity, probes and primers can be selected that comprise at least 20, 25, 30, 35, 40, 45, 50 or more consecutive nucleotides of a target nucleotide sequence.

Protein: A biological molecule, particularly a polypeptide, expressed by a gene and comprised of amino acids. A "polyprotein" is a protein that, after synthesis, is cleaved to produce several functionally distinct polypeptides.

Purified: The term "purified" does not require absolute purity; rather, it is intended as a relative term. Thus, for example, a purified protein preparation is one in which the subject protein is more pure than in its natural environment within a cell. Generally, a protein preparation is purified such that the protein represents at least 50% of the total protein content of the preparation.

Recombinant Nucleic Acid: A sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, for example, by genetic engineering techniques such as those described in Sambrook et al. (ed.), *Molecular Cloning: A Laboratory Manual*, 2nd ed., vol. 1-3, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid.

Sample: A portion, piece, or segment that is representative of a whole. This term encompasses any material, including for instance samples obtained from an animal, a plant, or the environment.

An "environmental sample" includes a sample obtained from inanimate objects or reservoirs within an indoor or outdoor environment. Environmental samples include, but are not limited to: soil, water, dust, and air samples; bulk samples, including building materials, furniture, and landfill contents; and other reservoir samples, such as animal refuse, harvested grains, and foodstuffs.

A "biological sample" is a sample obtained from a plant or animal subject. As used herein, biological samples include all samples useful for detection of viral infection in subjects, including, but not limited to: cells, tissues, and bodily fluids, such as blood; derivatives and fractions of blood (such as serum); extracted galls; biopsied or surgically removed tissue, including tissues that are, for example, unfixed, frozen, fixed in formalin and/or embedded in paraffin; tears; milk; skin scrapes; surface washings; urine; sputum; cerebrospinal fluid; prostate fluid; pus; bone marrow aspirates; BAL; saliva; cervical swabs; vaginal swabs; and oropharyngeal wash.

Sequence Identity The similarity between two nucleic acid sequences, or two amino acid sequences, is expressed in terms of the similarity between the sequences, otherwise referred to as sequence identity. Sequence identity is frequently measured in terms of percentage identity (or similarity or homology); the higher the percentage, the more similar the two sequences are.

Methods of alignment of sequences for comparison are well known in the art. Various programs and alignment algorithms are described in: Smith and Waterman (*Adv. Appl. Math.*, 2:482, 1981); Needleman and Wunsch (*J. Mol. Biol.*, 48:443, 1970); Pearson and Lipman (*Proc. Natl. Acad. Sci.*, 85:2444, 1988); Higgins and Sharp (*Gene*, 73:237-44, 1988); Higgins and Sharp (*CABIOS*, 5:151-53, 1989); Corpet et al. (*Nuc. Acids Res.*, 16:10881-90, 1988); Huang et al. (*Comp. Appls Biosci.*, 8:155-65, 1992); and Pearson et al. (*Meth. Mol. Biol.*, 24:307-31, 1994). Altschul et al. (*Nature Genet.*, 6:119-29, 1994) presents a detailed consideration of sequence alignment methods and homology calculations.

The alignment tools ALIGN (Myers and Miller, *CABIOS* 4:11-17, 1989) or LFASTA (Pearson and Lipman, 1988) may be used to perform sequence comparisons (Internet Program © 1996, W. R. Pearson and the University of Virginia, "fasta20u63" version 2.0u63, release date December 1996). ALIGN compares entire sequences against one another, while LFASTA compares regions of local similarity. These alignment tools and their respective tutorials are available on the Internet at the NCSA website. Alternatively, for comparisons of amino acid sequences of greater than about 30 amino acids, the "Blast 2 sequences" function can be employed using the default BLOSUM62 matrix set to default parameters, (gap existence cost of 11, and a per residue gap cost of 1). When aligning short peptides (fewer than around 30 amino acids), the alignment should be performed using the "Blast 2 sequences" function, employing the PAM30 matrix set to default parameters (open gap 9, extension gap 1 penalties). The BLAST sequence comparison system is available, for instance, from the NCBI web site; see also Altschul et al., *J. Mol. Biol.*, 215:403-10, 1990; Gish and States, *Nature Genet.*, 3:266-72, 1993; Madden et al., *Meth. Enzymol.*, 266:131-41, 1996; Altschul et al., *Nucleic Acids Res.*, 25:3389-402, 1997; and Zhang and Madden, *Genome Res.*, 7:649-56, 1997.

Orthologs (equivalent to proteins of other species) of proteins are in some instances characterized by possession of greater than 75% sequence identity counted over the full-length alignment with the amino acid sequence of specific protein using ALIGN set to default parameters. Proteins with even greater similarity to a reference sequence will show increasing percentage identities when assessed by this method, such as at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, or at least 98% sequence identity. In addition, sequence identity can be compared over the full length of one or both binding domains of the disclosed fusion proteins.

When significantly less than the entire sequence is being compared for sequence identity, homologous sequences will

typically possess at least 80% sequence identity over short windows of 10-20, and may possess sequence identities of at least 85%, at least 90%, at least 95%, or at least 99% depending on their similarity to the reference sequence. Sequence identity over such short windows can be determined using LFASTA; methods are described at the NCSA website. One of skill in the art will appreciate that these sequence identity ranges are provided for guidance only; it is entirely possible that strongly significant homologs could be obtained that fall outside of the ranges provided. Similar homology concepts apply for nucleic acids as are described for protein.

An alternative indication that two nucleic acid molecules are closely related is that the two molecules hybridize to each other under stringent conditions. Representative hybridization conditions are discussed above.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences, due to the degeneracy of the genetic code. It is understood that changes in nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that each encode substantially the same protein.

Specific Binding Agent: An agent that binds substantially only to a defined target. Thus a protein-specific binding agent binds substantially only the defined protein, or to a specific region within the protein. As used herein, a protein-specific binding agent includes antibodies and other agents that bind substantially to a specified polypeptide. The antibodies may be monoclonal or polyclonal antibodies that are specific for the polypeptide, as well as immunologically effective portions ("fragments") thereof.

The determination that a particular agent binds substantially only to a specific polypeptide may readily be made by using or adapting routine procedures. One suitable in vitro assay makes use of the Western blotting procedure (described in many standard texts, including Harlow and Lane, *Using Antibodies: A Laboratory Manual*, CSHL, New York, 1999).

Transformed: A "transformed" cell is a cell into which has been introduced a nucleic acid molecule by molecular biology techniques. The term encompasses all techniques by which a nucleic acid molecule might be introduced into such a cell, including transfection with viral vectors, transformation with plasmid vectors, and introduction of naked DNA by electroporation, lipofection, and particle gun acceleration.

Vector: A nucleic acid molecule as introduced into a host cell, thereby producing a transformed host cell. A vector may include nucleic acid sequences that permit it to replicate in a host cell, such as an origin of replication. A vector may also include one or more selectable marker genes and other genetic elements known in the art.

Virus: Microscopic infectious organism that reproduces inside living cells. A virus typically consists essentially of a core of a single nucleic acid surrounded by a protein coat, and has the ability to replicate only inside a living cell. "Viral replication" is the production of additional virus by the occurrence of at least one viral life cycle. A virus may subvert the host cells' normal functions, causing the cell to behave in a manner determined by the virus. For example, a viral infection may result in a cell producing a cytokine, or responding to a cytokine, when the uninfected cell does not normally do so.

"Coronaviruses" are large, enveloped, RNA viruses that cause respiratory and enteric diseases in humans and other animals. Coronavirus genomes are non-segmented, single-stranded, positive-sense RNA, approximately 27-31 kb in length. Genomes have a 5' methylated cap and 3' poly-A tail, and function directly as mRNA. Host cell entry occurs via endocytosis and membrane fusion, and replication occurs in

the cytoplasm. Initially, the 5' 20 kb of the positive-sense genome is translated to produce a viral polymerase, which then produces a full-length negative-sense strand used as a template to produce subgenomic mRNA as a "nested set" of transcripts. Assembly occurs by budding into the golgi apparatus, and particles are transported to the surface of the cell and released.

III. Overview of Several Embodiments

A newly isolated human coronavirus (SARS-CoV) is disclosed herein. The entire genomic nucleic acid sequence of this virus is also provided herein. Also disclosed are the nucleic acid sequences of the SARS-CoV ORFs, and the polypeptide sequences encoded by these ORFs. Pharmaceutical and immune stimulatory compositions are also disclosed that include one or more SARS-CoV viral nucleic acids, polypeptides encoded by these viral nucleic acids and antibodies that bind to a SARS-CoV polypeptide or SARS-CoV polypeptide fragment.

In one embodiment, a method is provided for detecting the presence of SARS-CoV in a sample. This method includes contacting the sample with a pair of nucleic acid primers that hybridize to a SARS-CoV nucleic acid, wherein at least one primer is 5'-end labeled with a reporter dye, amplifying the SARS-CoV nucleic acid or a fragment thereof from the sample utilizing the pair of nucleic acid primers, electrophoresing the amplified products, and detecting the 5'-end labeled reporter dye, thereby detecting a SARS-CoV. In a specific, non-limiting example, the amplification utilizes RT-PCR. In a further specific example of the provided method, at least one of the nucleic acid primers that hybridize to a SARS-CoV nucleic acid includes a sequence as set forth in any one of SEQ ID NOS: 13-15.

In another example of the provided method, detecting the presence of SARS-CoV in a sample includes contacting the sample with a pair of nucleic acid primers that hybridize to a SARS-CoV nucleic acid, amplifying the SARS-CoV nucleic acid or a fragment thereof from the sample utilizing the pair of nucleic acid primers, adding to the amplified SARS-CoV nucleic acid or the fragment thereof a TaqMan SARS-CoV probe that hybridizes to the SARS-CoV nucleic acid, wherein the TaqMan SARS-CoV probe is labeled with a 5'-reporter dye and a 3'-quencher dye, performing one or more additional rounds of amplification, and detecting fluorescence of the 5'-reporter dye, thereby detecting a SARS-CoV. In a specific, non-limiting example, the amplification utilizes RT-PCR. In a further specific example of the provided method, at least one of the nucleic acid primers that hybridize to a SARS-CoV nucleic acid and/or the TaqMan SARS-CoV probe that hybridizes to the SARS-CoV nucleic acid includes a sequence as set forth in any one of SEQ ID NOS: 16-33.

In another embodiment, a method is provided for detecting a SARS-CoV in a biological sample that contains antibodies. This method includes contacting the biological sample with a SARS-CoV-specific antigen, wherein the antigen includes a SARS-CoV organism and determining whether a binding reaction occurs between the SARS-CoV-specific antigen and an antibody in the biological sample if such is present, thereby detecting SARS-CoV.

In a further embodiment, a method is provided for detecting a SARS-CoV in a biological sample that contains polypeptides and/or fragments thereof. This method includes contacting the biological sample with a SARS-CoV-specific antibody and determining whether a binding reaction occurs between the SARS-CoV-specific antibody and a SARS-CoV polypeptide or fragment thereof in the biological sample if

such is present, thereby detecting SARS-CoV. In a specific, non-limiting example, determining whether a binding reaction occurs between the SARS-CoV-specific antibody and a SARS-CoV polypeptide or fragment thereof is carried out in situ or in a tissue sample. In a further specific example, determining whether a binding reaction occurs between the SARS-CoV-specific antibody and a SARS-CoV polypeptide or fragment thereof includes an immunohistochemical assay.

An additional embodiment includes a kit for detecting a SARS-CoV in a sample, including a pair of nucleic acid primers that hybridize under stringent conditions to a SARS-CoV nucleic acid, wherein one primer is 5'-end labeled with a reporter dye. In a specific, non-limiting example, at least one of the nucleic acid primers that hybridize to a SARS-CoV nucleic acid includes a sequence as set forth in any one of SEQ ID NOs: 13-15.

Another example of the provided kit includes a pair of nucleic acid primers that hybridize under high stringency conditions to a SARS-CoV nucleic acid and a TaqMan SARS-CoV probe that hybridizes to the SARS-CoV nucleic acid, wherein the TaqMan SARS-CoV probe is labeled with a 5'-reporter dye and a 3'-quencher dye. In a specific, non-limiting example, at least one of the nucleic acid primers that hybridize to a SARS-CoV nucleic acid and/or the TaqMan SARS-CoV probe that hybridizes to the SARS-CoV nucleic acid includes a sequence as set forth in any one of SEQ ID NOs: 16-33.

Also disclosed herein is a composition including an isolated SARS-CoV organism. In one embodiment, the isolated SARS-CoV organism is an inactive isolated SARS-CoV organism. In another embodiment, the composition includes at least one component selected from the group consisting of pharmaceutically acceptable carriers, adjuvants and combinations of two or more thereof. In yet another embodiment, the composition is introduced into a subject, thereby eliciting an immune response against a SARS-CoV antigenic epitope in a subject.

IV. SARS-CoV Nucleotide and Amino Acid Sequences

The current disclosure provides an isolated SARS-CoV genome, isolated SARS-CoV polypeptides, and isolated nucleic acid molecules encoding the same. In one embodiment, the isolated SARS-CoV genome has a sequence as shown in SEQ ID NO: 1 or an equivalent thereof. Polynucleotides encoding a SARS-CoV polypeptide (encoded by an ORF from within the genome) are also provided, and are termed SARS-CoV nucleic acid molecules. These nucleic acid molecules include DNA, cDNA and RNA sequences which encode a SARS-CoV polypeptide. Specific, non-limiting examples of a SARS-CoV nucleic acid molecule encoding an ORF are nucleic acid 265 to nucleic acid 13,398 of SEQ ID NO: 1 (encoding SARS-CoV 1a, SEQ ID NO: 2), nucleic acid 13,398 to 21,482 of SEQ ID NO: 1 (encoding SARS-CoV 1b, SEQ ID NO: 3), nucleic acid 21,492 to 25,256 of SEQ ID NO: 1 (encoding SARS-CoV S, SEQ ID NO: 4), nucleic acid 25,268 to 26,089 of SEQ ID NO: 1 (encoding SARS-CoV X1, SEQ ID NO: 5), nucleic acid 25,689 to 26,150 of SEQ ID NO: 1 (encoding SARS-CoV X2, SEQ ID NO: 6), nucleic acid 26,117 to 26,344 of SEQ ID NO: 1 (encoding SARS-CoV E, SEQ ID NO: 7), nucleic acid 26,398 to 27,060 of SEQ ID NO: 1 (encoding SARS-CoV M, SEQ ID NO: 8), nucleic acid 27,074 to 27,262 of SEQ ID NO: 1 (encoding SARS-CoV X3, SEQ ID NO: 9), nucleic acid 27,273 to 27,638 of SEQ ID NO: 1 (encoding SARS-CoV X4, SEQ ID NO: 10), nucleic acid 27,864 to 28,115 of SEQ ID

NO: 1 (encoding SARS-CoV X5, SEQ ID NO: 11), and nucleic acid 28,120 to 29,385 of SEQ ID NO: 1 (encoding SARS-CoV N, SEQ ID NO: 12).

Oligonucleotide primers and probes derived from the SARS-CoV genome (SEQ ID NO: 1) are also encompassed within the scope of the present disclosure. Such oligonucleotide primers and probes may comprise a sequence of at least about 15 consecutive nucleotides of the SARS-CoV nucleic acid sequence, such as at least about 20, 25, 30, 35, 40, 45, or 50 or more consecutive nucleotides. These primers and probes may be obtained from any region of the disclosed SARS-CoV genome (SEQ ID NO: 1), including particularly from any of the ORFs disclosed herein. Specific, non-limiting examples of oligonucleotide primers derived from the SARS-CoV genome (SEQ ID NO: 1) include: Cor-p-F2 (SEQ ID NO: 13), Cor-p-F3 (SEQ ID NO: 14), Cor-p-R1 (SEQ ID NO: 15), SARS1-F (SEQ ID NO: 16), SARS1-R (SEQ ID NO: 17), SARS2-F (SEQ ID NO: 19), SARS2-R (SEQ ID NO: 20), SARS3-F (SEQ ID NO: 22), SARS3-R (SEQ ID NO: 23), N3-F (SEQ ID NO: 25), N3-R (SEQ ID NO: 26), 3'NTR-F (SEQ ID NO: 28), 3'NTR-R (SEQ ID NO: 29), M-F (SEQ ID NO: 31), and M-R (SEQ ID NO: 32). Specific, non-limiting examples of oligonucleotide probes derived from the SARS-CoV genome (SEQ ID NO: 1) include: SARS1-P (SEQ ID NO: 18), SARS2-P (SEQ ID NO: 21), SARS3-P (SEQ ID NO: 24), N3-P (SEQ ID NO: 27), 3'NTR-P (SEQ ID NO: 30), and M-P (SEQ ID NO: 33).

Nucleic acid molecules encoding a SARS-CoV polypeptide can be operatively linked to regulatory sequences or elements. Regulatory sequences or elements include, but are not limited to promoters, enhancers, transcription terminators, a start codon (for example, ATG), stop codons, and the like.

Additionally, nucleic acid molecules encoding a SARS-CoV polypeptide can be inserted into an expression vector. Specific, non-limiting examples of vectors include, plasmids, bacteriophages, cosmids, animal viruses and yeast artificial chromosomes (YACs) (Burke et al., *Science* 236:806-12, 1987). Such vectors may then be introduced into a variety of hosts including somatic cells, and simple or complex organisms, such as bacteria, fungi (Timberlake and Marshall, *Science* 244:1313-17, 1989), invertebrates, plants (Gasser et al., *Plant Cell* 1:15-24, 1989), and animals (Pursel et al., *Science* 244:1281-88, 1989).

Transformation of a host cell with an expression vector carrying a nucleic acid molecule encoding a SARS-CoV polypeptide may be carried out by conventional techniques, as are well known to those skilled in the art. By way of example, where the host is prokaryotic, such as *E. coli*, competent cells that are capable of DNA uptake can be prepared from cells harvested after exponential growth phase and subsequently treated by the CaCl₂ method using procedures well known in the art. Alternatively, MgCl₂ or RbCl can be used. Transformation can also be performed after forming a protoplast of the host cell if desired, or by electroporation.

When the host is a eukaryote, methods of transfection of DNA, such as calcium phosphate coprecipitates, conventional mechanical procedures such as microinjection, electroporation, insertion of a plasmid encased in liposomes, or virus vectors, may be used. Eukaryotic cells can also be cotransformed with SARS-CoV nucleic acid molecules, and a second foreign DNA molecule encoding a selectable phenotype, such as the herpes simplex thymidine kinase gene. Another method is to use a eukaryotic viral vector, such as simian virus 40 or bovine papilloma virus, to transiently infect or transform eukaryotic cells and express the protein

(see, for example, *Eukaryotic Viral Vectors*, Cold Spring Harbor Laboratory, Gluzman ed., 1982).

The SARS-CoV polypeptides of this disclosure include proteins encoded by any of the ORFs disclosed herein, and equivalents thereof. Specific, non-limiting examples of SARS-CoV proteins are provided in SEQ ID NOs: 2-12. Fusion proteins are also contemplated that include a heterologous amino acid sequence chemically linked to a SARS-CoV polypeptide. Exemplary heterologous sequences include short amino acid sequence tags (such as six histidine residues), as well as a fusion of other proteins (such as c-myc or green fluorescent protein fusions). Epitopes of the SARS-CoV proteins, that are recognized by an antibody or that bind the major histocompatibility complex, and can be used to induce a SARS-CoV-specific immune response, are also encompassed by this disclosure.

Methods for expressing large amounts of protein from a cloned gene introduced into *E. coli* may be utilized for the purification and functional analysis of proteins. For example, fusion proteins consisting of amino terminal peptides encoded by a portion of the *E. coli* lacZ or trpE gene linked to SARS-CoV proteins may be used to prepare polyclonal and monoclonal antibodies against these proteins.

Intact native protein may also be produced in *E. coli* in large amounts for functional studies. Methods and plasmid vectors for producing fusion proteins and intact native proteins in bacteria are described by Sambrook et al. (ed.), *Molecular Cloning: A Laboratory Manual*, 2nd ed., vol. 1-3, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989. Such fusion proteins may be made in large amounts, are easy to purify, and can be used to elicit antibody response. Native proteins can be produced in bacteria by placing a strong, regulated promoter and an efficient ribosome-binding site upstream of the cloned gene. If low levels of protein are produced, additional steps may be taken to increase protein production; if high levels of protein are produced, purification is relatively easy. Suitable methods are presented by Sambrook et al. (ed.), *Molecular Cloning: A Laboratory Manual*, 2nd ed., vol. 1-3, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989, and are well known in the art. Often, proteins expressed at high levels are found in insoluble inclusion bodies. Methods for extracting proteins from these aggregates are described by Sambrook et al. (ed.), *Molecular Cloning: A Laboratory Manual*, 2nd ed., vol. 1-3, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989.

Isolation and purification of recombinantly expressed proteins may be carried out by conventional means including preparative chromatography and immunological separations. Additionally, the proteins can be chemically synthesized by any of a number of manual or automated methods of synthesis known in the art.

V. Specific Binding Agents

The disclosure provides specific binding agents that bind to SARS-CoV polypeptides disclosed herein. The binding agent may be useful for purifying and detecting the polypeptides, as well as for detection and diagnosis of SARS-CoV. Examples of the binding agents are a polyclonal or monoclonal antibody, and fragments thereof, that bind to any of the SARS-CoV polypeptides disclosed herein.

Monoclonal or polyclonal antibodies may be raised to recognize a SARS-CoV polypeptide described herein, or a fragment or variant thereof. Optimally, antibodies raised against these polypeptides would specifically detect the polypeptide with which the antibodies are generated. That is, antibodies

raised against a specific SARS-CoV polypeptide will recognize and bind that polypeptide, and will not substantially recognize or bind to other polypeptides or antigens. The determination that an antibody specifically binds to a target polypeptide is made by any one of a number of standard immunoassay methods; for instance, the Western blotting technique (Sambrook et al. (ed.), *Molecular Cloning: A Laboratory Manual*, 2nd ed., vol. 1-3, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989).

Substantially pure SARS-CoV recombinant polypeptide antigens suitable for use as immunogen may be isolated from the transformed cells described above, using methods well known in the art. Monoclonal or polyclonal antibodies to the antigens may then be prepared.

Monoclonal antibodies to the polypeptides can be prepared from murine hybridomas according to the classic method of Kohler & Milstein (*Nature* 256:495-97, 1975), or a derivative method thereof. Briefly, a mouse is repetitively inoculated with a few micrograms of the selected protein immunogen (for example, a polypeptide comprising at least one SARS-CoV-specific epitope, a portion of a polypeptide comprising at least one SARS-CoV-specific epitope, or a synthetic peptide comprising at least one SARS-CoV-specific epitope) over a period of a few weeks. The mouse is then sacrificed, and the antibody-producing cells of the spleen isolated. The spleen cells are fused by means of polyethylene glycol with mouse myeloma cells, and the excess unfused cells destroyed by growth of the system on selective media comprising aminopterin (HAT media). The successfully fused cells are diluted and aliquots of the dilution placed in wells of a microtiter plate where growth of the culture is continued. Antibody-producing clones are identified by detection of antibody in the supernatant fluid of the wells by immunoassay procedures, such as ELISA, as originally described by Engvall (*Meth. Enzymol.*, 70:419-39, 1980), or a derivative method thereof. Selected positive clones can be expanded and their monoclonal antibody product harvested for use. Detailed procedures for monoclonal antibody production are described in Harlow and Lane, *Using Antibodies: A Laboratory Manual*, CSHL, New York, 1999.

Polyclonal antiserum containing antibodies can be prepared by immunizing suitable animals with a polypeptide comprising at least one SARS-CoV-specific epitope, a portion of a polypeptide comprising at least one SARS-CoV-specific epitope, or a synthetic peptide comprising at least one SARS-CoV-specific epitope, which can be unmodified or modified, to enhance immunogenicity.

Effective antibody production (whether monoclonal or polyclonal) is affected by many factors related both to the antigen and the host species. For example, small molecules tend to be less immunogenic than others and may require the use of carriers and adjuvant. Also, host animals vary in response to site of inoculations and dose, with either inadequate or excessive doses of antigen resulting in low titer antisera. Small doses (ng level) of antigen administered at multiple intradermal sites appear to be most reliable. An effective immunization protocol for rabbits can be found in Vaitukaitis et al. (*J. Clin. Endocrinol. Metab.*, 33:988-91, 1971).

Booster injections can be given at regular intervals, and antiserum harvested when the antibody titer thereof, as determined semi-quantitatively, for example, by double immunodiffusion in agar against known concentrations of the antigen, begins to fall. See, for example, Ouchterlony et al., *Handbook of Experimental Immunology*, Wier, D. (ed.), Chapter 19, Blackwell, 1973. A plateau concentration of antibody is usually in the range of 0.1 to 0.2 mg/ml of serum

(about 12 μ M). Affinity of the antisera for the antigen is determined by preparing competitive binding curves, as described, for example, by Fisher (*Manual of Clinical Immunology*, Ch. 42, 1980).

Antibody fragments may be used in place of whole antibodies and may be readily expressed in prokaryotic host cells. Methods of making and using immunologically effective portions of monoclonal antibodies, also referred to as "antibody fragments," are well known and include those described in Better & Horowitz, *Methods Enzymol.* 178:476-96, 1989; Glockshuber et al., *Biochemistry* 29:1362-67, 1990; and U.S. Pat. Nos. 5,648,237 (Expression of Functional Antibody Fragments); 4,946,778 (Single Polypeptide Chain Binding Molecules); and 5,455,030 (Immunotherapy Using Single Chain Polypeptide Binding Molecules), and references cited therein. Conditions whereby a polypeptide/binding agent complex can form, as well as assays for the detection of the formation of a polypeptide/binding agent complex and quantitation of binding affinities of the binding agent and polypeptide, are standard in the art. Such assays can include, but are not limited to, Western blotting, immunoprecipitation, immunofluorescence, immunocytochemistry, immunohistochemistry, fluorescence activated cell sorting (FACS), fluorescence in situ hybridization (FISH), immunomagnetic assays, ELISA, ELISPOT (Coligan et al., *Current Protocols in Immunology*, Wiley, NY, 1995), agglutination assays, flocculation assays, cell panning, and the like, as are well known to one of skill in the art.

Binding agents of this disclosure can be bound to a substrate (for example, beads, tubes, slides, plates, nitrocellulose sheets, and the like) or conjugated with a detectable moiety, or both bound and conjugated. The detectable moieties contemplated for the present disclosure can include, but are not limited to, an immunofluorescent moiety (for example, fluorescein, rhodamine), a radioactive moiety (for example, 32 P, 125 I, 35 S), an enzyme moiety (for example, horseradish peroxidase, alkaline phosphatase), a colloidal gold moiety, and a biotin moiety. Such conjugation techniques are standard in the art (for example, see Harlow and Lane, *Using Antibodies: A Laboratory Manual*, CSHL, New York, 1999; Yang et al., *Nature*, 382:319-24, 1996).

VI. Detection and Diagnosis of SARS-CoV

A. Nucleic Acid Based Methods of Detection and Diagnosis

A major application of the SARS-CoV sequence information presented herein is in the area of detection and diagnostic testing for SARS-CoV infection. Methods for screening a subject to determine if the subject has been or is currently infected with SARS-CoV are disclosed herein.

One such method includes providing a sample, which sample includes a nucleic acid such as DNA or RNA, and providing an assay for detecting in the sample the presence of a SARS-CoV nucleic acid molecule. Suitable samples include all biological samples useful for detection of viral infection in subjects, including, but not limited to, cells, tissues (for example, lung and kidney), bodily fluids (for example, blood, serum, urine, saliva, sputum, and cerebrospinal fluid), bone marrow aspirates, BAL, and oropharyngeal wash. Additional suitable samples include all environmental samples useful for detection of viral presence in the environment, including, but not limited to, a sample obtained from inanimate objects or reservoirs within an indoor or outdoor environment. The detection in the sample of a SARS-CoV nucleic acid molecule may be performed by a number of methodologies, non-limiting examples of which are outlined below.

In one embodiment, detecting in the sample the presence of a SARS-CoV nucleic acid molecule includes the amplification of a SARS-CoV nucleic acid sequence (or a fragment thereof). Any nucleic acid amplification method can be used.

In one specific, non-limiting example, PCR is used to amplify the SARS-CoV nucleic acid sequence(s). In another non-limiting example, RT-PCR can be used to amplify the SARS-CoV nucleic acid sequences. In an additional non-limiting example, transcription-mediated amplification can be used to amplify the SARS-CoV nucleic acid sequences.

In some embodiments, a pair of SARS-CoV-specific primers are utilized in the amplification reaction. One or both of the primers can be end-labeled (for example, radiolabeled, fluoresceinated, or biotinylated). In one specific, non-limiting example, at least one of the primers is 5'-end labeled with the reporter dye 6-carboxyfluorescein (6-FAM). The pair of primers includes an upstream primer (which binds 5' to the downstream primer) and a downstream primer (which binds 3' to the upstream primer). In one embodiment, either the upstream primer or the downstream primer is labeled. Specific, non-limiting examples of SARS-CoV-specific primers include, but are not limited to: Cor-p-F2 (SEQ ID NO: 13), Cor-p-F3 (SEQ ID NO: 14), Cor-p-R1 (SEQ ID NO: 15), SARS1-F (SEQ ID NO: 16), SARS1-R (SEQ ID NO: 17), SARS2-F (SEQ ID NO: 19), SARS2-R (SEQ ID NO: 20), SARS3-F (SEQ ID NO: 22), SARS3R (SEQ ID NO: 23), N3-F (SEQ ID NO: 25), N3-R (SEQ ID NO: 26), 3'NTR-F (SEQ ID NO: 28), 3'NTR-R (SEQ ID NO: 29), M-F (SEQ ID NO: 31), and M-R (SEQ ID NO: 32). Additional primer pairs can be generated, for instance, to amplify any of the specific ORFs described herein, using well known primer design principles and methods.

In one specific, non-limiting example, electrophoresis is used to detect amplified SARS-CoV-specific sequences. Electrophoresis can be automated using many methods well known in the art. In one embodiment, a genetic analyzer is used, such as an ABI 3100 Prism Genetic Analyzer (PE Applied Biosystems, Foster City, Calif.), wherein the bands are analyzed using GeneScan software (PE Applied Biosystems, Foster City, Calif.).

In another specific, non-limiting example, hybridization assays are used to detect amplified SARS-CoV-specific sequences using distinguishing oligonucleotide probes. Such probes include "TaqMan" probes. TaqMan probes consist of an oligonucleotide with a reporter at the 5'-end and a quencher at the 3'-end. In one specific, non-limiting example, the reporter is 6-FAM and the quencher is Blackhole Quencher (Biosearch Tech., Inc., Novato, Calif.). When the probe is intact, the proximity of the reporter to the quencher results in suppression of reporter fluorescence, primarily by fluorescence resonance energy transfer. If the target of interest is present, the TaqMan probe specifically hybridizes between the forward and reverse primer sites during the PCR annealing step. In the process of PCR elongation, the 5'-3' nucleolytic activity of the Taq DNA polymerase cleaves the hybridized probe between the reporter and the quencher. The probe fragments are then displaced from the target, and polymerization of the strand continues. Taq DNA polymerase does not cleave non-hybridized probe, and cleaves the hybridized probe only during polymerization. The 3'-end of the probe is blocked to prevent extension of the probe during PCR. The 5'-3' nuclease cleavage of the hybridized probe occurs in every cycle and does not interfere with the exponential accumulation of PCR product. Accumulation of PCR products is detected directly by monitoring the increase in fluorescence of the released reporter. The increase in fluorescence signal is detected only if the target sequence is complementary to the

probe and is amplified during PCR. Therefore, non-specific amplification is not detected. SARS-CoV-specific TaqMan probes of the present disclosure include, but are not limited to: SARS1-P (SEQ ID NO: 18), SARS2-P (SEQ ID NO: 21), SARS3-P (SEQ ID NO: 24), N3-P (SEQ ID NO: 27), 3'NTR-P (SEQ ID NO: 30), and M-P (SEQ ID NO: 33), and hybridization assays include, but are not limited to, a real-time RT-PCR assay.

B. Protein Based Methods of Detection and Diagnosis

The present disclosure further provides methods of detecting a SARS-CoV antigen in a sample, and/or diagnosing SARS-CoV infection in a subject by detecting a SARS-CoV antigen. Examples of such methods comprise contacting the sample with a SARS-CoV-specific binding agent under conditions whereby an antigen/binding agent complex can form; and detecting formation of the complex, thereby detecting SARS-CoV antigen in a sample and/or diagnosing SARS-CoV infection in a subject. It is contemplated that at least certain antigens will be on an intact SARS-CoV virion, will be a SARS-CoV-encoded protein displayed on the surface of a SARS-CoV-infected cell expressing the antigen, or will be a fragment of the antigen. Contemplated samples subject to analysis by these methods can comprise any sample, such as a clinical sample, useful for detection of viral infection in a subject.

Methods for detecting antigens in a sample are discussed, for example, in Ausubel et al. *Short Protocols in Molecular Biology*, 4th ed., John Wiley & Sons, Inc., 1999. Enzyme immunoassays such as IFA, ELISA and immunoblotting can be readily adapted to accomplish the detection of SARS-CoV antigens according to the methods of this disclosure. An ELISA method effective for the detection of soluble SARS-CoV antigens is the direct competitive ELISA. This method is most useful when a specific SARS-CoV antibody and purified SARS-CoV antigen are available. Briefly: 1) coat a substrate (for example, a microtiter plate) with a sample suspected of containing a SARS-CoV antigen; 2) contact the bound SARS-CoV antigen with a SARS-CoV-specific antibody bound to a detectable moiety (for example, horseradish peroxidase enzyme or alkaline phosphatase enzyme); 3) add purified inhibitor SARS-CoV antigen; 4) contact the above with the substrate for the enzyme; and 5) observe/measure inhibition of color change or fluorescence and quantitate antigen concentration (for example, using a microtiter plate reader).

An additional ELISA method effective for the detection of soluble SARS-CoV antigens is the antibody-sandwich ELISA. This method is frequently more sensitive in detecting antigen than the direct competitive ELISA method. Briefly: 1) coat a substrate (for example, a microtiter plate) with a SARS-CoV-specific antibody; 2) contact the bound SARS-CoV antibody with a sample suspected of containing a SARS-CoV antigen; 3) contact the above with SARS-CoV-specific antibody bound to a detectable moiety (for example, horseradish peroxidase enzyme or alkaline phosphatase enzyme); 4) contact the above with the substrate for the enzyme; and 5) observe/measure color change or fluorescence and quantitate antigen concentration (for example, using a microtiter plate reader).

An ELISA method effective for the detection of cell-surface SARS-CoV antigens is the direct cellular ELISA. Briefly, cells suspected of exhibiting a cell-surface SARS-CoV antigen are fixed (for example, using glutaraldehyde) and incubated with a SARS-CoV-specific antibody bound to a detectable moiety (for example, horseradish peroxidase enzyme or alkaline phosphatase enzyme). Following a wash

to remove unbound antibody, substrate for the enzyme is added and color change or fluorescence is observed/measured.

The present disclosure further provides methods of detecting a SARS-CoV-reactive antibody in a sample, and/or diagnosing SARS-CoV infection in a subject by detecting a SARS-CoV-reactive antibody. Examples of such methods comprise contacting the sample with a SARS-CoV polypeptide of this disclosure under conditions whereby a polypeptide/antibody complex can form; and detecting formation of the complex, thereby detecting SARS-CoV antibody in a sample and/or diagnosing SARS-CoV infection in a subject. Contemplated samples subject to analysis by these methods can comprise any sample, such as a clinical sample, as described herein as being useful for detection of viral infection in a subject.

Methods for detecting antibodies in a sample are discussed, for example, in Ausubel et al. *Short Protocols in Molecular Biology*, 4th ed., John Wiley & Sons, Inc., 1999. Enzyme immunoassays such as IFA, ELISA and immunoblotting can be readily adapted to accomplish the detection of SARS-CoV antibodies according to the methods of this disclosure. An ELISA method effective for the detection of specific SARS-CoV antibodies is the indirect ELISA method. Briefly: 1) bind a SARS-CoV polypeptide to a substrate (for example, a microtiter plate); 2) contact the bound polypeptide with a sample suspected of containing SARS-CoV antibody; 3) contact the above with a secondary antibody bound to a detectable moiety which is reactive with the bound antibody (for example, horseradish peroxidase enzyme or alkaline phosphatase enzyme); 4) contact the above with the substrate for the enzyme; and 5) observe/measure color change or fluorescence.

Another immunologic technique that can be useful in the detection of SARS-CoV antibodies uses monoclonal antibodies for detection of antibodies specifically reactive with SARS-CoV polypeptides in a competitive inhibition assay. Briefly, a sample suspected of containing SARS-CoV antibodies is contacted with a SARS-CoV polypeptide of this disclosure which is bound to a substrate (for example, a microtiter plate). Excess sample is thoroughly washed away. A labeled (for example, enzyme-linked, fluorescent, radioactive, and the like) monoclonal antibody specific for the SARS-CoV polypeptide is then contacted with any previously formed polypeptide-antibody complexes and the amount of monoclonal antibody binding is measured. The amount of inhibition of monoclonal antibody binding is measured relative to a control (no monoclonal antibody), allowing for detection and measurement of antibody in the sample. The degree of monoclonal antibody inhibition can be a very specific assay for detecting SARS-CoV. Monoclonal antibodies can also be used for direct detection of SARS-CoV in cells or tissue samples by, for example, IFA analysis according to standard methods.

As a further example, a micro-agglutination test can be used to detect the presence of SARS-CoV antibodies in a sample. Briefly, latex beads, red blood cells or other agglutinable particles are coated with a SARS-CoV polypeptide of this disclosure and mixed with a sample, such that antibodies in the sample that are specifically reactive with the antigen crosslink with the antigen, causing agglutination. The agglutinated polypeptide-antibody complexes form a precipitate, visible with the naked eye or measurable by spectrophotometer. In a modification of the above test, SARS-CoV-specific antibodies of this disclosure can be bound to the agglutinable particles and SARS-CoV antigen in the sample thereby detected.

VII. Pharmaceutical and Immune Stimulatory Compositions and Uses Thereof

Pharmaceutical compositions including SARS-CoV nucleic acid sequences, SARS-CoV polypeptides, or antibodies that bind these polypeptides, are also encompassed by the present disclosure. These pharmaceutical compositions include a therapeutically effective amount of one or more SARS-CoV polypeptides, one or more nucleic acid molecules encoding a SARS-CoV polypeptide, or an antibody that binds a SARS-CoV polypeptide, in conjunction with a pharmaceutically acceptable carrier.

Disclosed herein are substances suitable for use as immune stimulatory compositions for the inhibition or treatment of SARS. Particular immune stimulatory compositions are directed against SARS-CoV, and include antigens obtained from SARS-CoV. In one embodiment, an immune stimulatory composition contains attenuated SARS-CoV. Methods of viral attenuation are well known in the art, and include, but are not limited to, high serial passage (for example, in susceptible host cells under specific environmental conditions to select for attenuated virions), exposure to a mutagenic agent (for example, a chemical mutagen or radiation), genetic engineering using recombinant DNA technology (for example, using gene replacement or gene knockout to disable one or more viral genes), or some combination thereof.

In another embodiment, the immune stimulatory composition contains inactivated SARS-CoV. Methods of viral inactivation are well known in the art, and include, but are not limited to, heat and chemicals (for example, formalin, β -propiolactone, and ethylenimines).

In yet another embodiment, the immune stimulatory composition contains a nucleic acid vector that includes SARS-CoV nucleic acid molecules described herein, or that includes a nucleic acid sequence encoding an immunogenic polypeptide or polypeptide fragment of SARS-CoV or derived from SARS-CoV, such as a polypeptide that encodes a surface protein of SARS-CoV.

In a further embodiment, the immune stimulatory composition contains a SARS-CoV subunit, such as glycoprotein, major capsid protein, or other gene products found to elicit humoral and/or cell mediated immune responses.

The provided immune stimulatory SARS-CoV polypeptides, constructs or vectors encoding such polypeptides, are combined with a pharmaceutically acceptable carrier or vehicle for administration as an immune stimulatory composition to human or animal subjects. In some embodiments, more than one immune stimulatory SARS-CoV polypeptide may be combined to form a single preparation.

The immunogenic formulations may be conveniently presented in unit dosage form and prepared using conventional pharmaceutical techniques. Such techniques include the step of bringing into association the active ingredient and the pharmaceutical carrier(s) or excipient(s). In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers. Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of a sterile liquid carrier, for example, water for injections, immediately prior to use.

Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets commonly used by one of ordinary skill in the art.

In certain embodiments, unit dosage formulations are those containing a dose or unit, or an appropriate fraction thereof, of the administered ingredient. It should be understood that in addition to the ingredients particularly mentioned above, formulations encompassed herein may include other agents commonly used by one of ordinary skill in the art.

The compositions provided herein, including those for use as immune stimulatory compositions, may be administered through different routes, such as oral, including buccal and sublingual, rectal, parenteral, aerosol, nasal, intramuscular, subcutaneous, intradermal, and topical. They may be administered in different forms, including but not limited to solutions, emulsions and suspensions, microspheres, particles, microparticles, nanoparticles, and liposomes.

The volume of administration will vary depending on the route of administration. By way of example, intramuscular injections may range from about 0.1 ml to about 1.0 ml. Those of ordinary skill in the art will know appropriate volumes for different routes of administration.

A relatively recent development in the field of immune stimulatory compounds (for example, vaccines) is the direct injection of nucleic acid molecules encoding peptide antigens (broadly described in Janeway & Travers, *Immunobiology: The Immune System In Health and Disease*, page 13.25, Garland Publishing, Inc., New York, 1997; and McDonnell & Askari, *N. Engl. J. Med.* 334:42-45, 1996). Vectors that include nucleic acid molecules described herein, or that include a nucleic acid sequence encoding an immunogenic SARS-CoV polypeptide may be utilized in such DNA vaccination methods.

Thus, the term "immune stimulatory composition" as used herein also includes nucleic acid vaccines in which a nucleic acid molecule encoding a SARS-CoV polypeptide is administered to a subject in a pharmaceutical composition. For genetic immunization, suitable delivery methods known to those skilled in the art include direct injection of plasmid DNA into muscles (Wolff et al., *Hum. Mol. Genet.* 1:363, 1992), delivery of DNA complexed with specific protein carriers (Wu et al., *J. Biol. Chem.* 264:16985, 1989), co-precipitation of DNA with calcium phosphate (Benvenisty and Reshef, *Proc. Natl. Acad. Sci.* 83:9551, 1986), encapsulation of DNA in liposomes (Kaneda et al., *Science* 243:375, 1989), particle bombardment (Tang et al., *Nature* 356:152, 1992; Eisenbraun et al., *DNA Cell Biol.* 12:791, 1993), and in vivo infection using cloned retroviral vectors (Seeger et al., *Proc. Natl. Acad. Sci.* 81:5849, 1984). Similarly, nucleic acid vaccine preparations can be administered via viral carrier.

The amount of immunostimulatory compound in each dose of an immune stimulatory composition is selected as an amount that induces an immunostimulatory or immunoprotective response without significant, adverse side effects. Such amount will vary depending upon which specific immunogen is employed and how it is presented. Initial injections may range from about 1 μ g to about 1 mg, with some embodiments having a range of about 10 μ g to about 800 μ g, and still other embodiments a range of from about 25 μ g to about 500 μ g. Following an initial administration of the immune stimulatory composition, subjects may receive one or several booster administrations, adequately spaced. Booster administrations may range from about 1 μ g to about 1 mg, with other embodiments having a range of about 10 μ g to about 750 μ g, and still others a range of about 50 μ g to about 500 μ g.

Periodic boosters at intervals of 1-5 years, for instance three years, may be desirable to maintain the desired levels of protective immunity.

It is also contemplated that the provided immunostimulatory molecules and compositions can be administered to a subject indirectly, by first stimulating a cell in vitro, which stimulated cell is thereafter administered to the subject to elicit an immune response. Additionally, the pharmaceutical or immune stimulatory compositions or methods of treatment may be administered in combination with other therapeutic treatments.

VIII. Kits

Also provided herein are kits useful in the detection and/or diagnosis of SARS-CoV. This includes kits for use with nucleic acid and protein detection methods, such as those disclosed herein.

The SARS-CoV-specific oligonucleotide primers and probes described herein can be supplied in the form of a kit for use in detection of SARS-CoV. In such a kit, an appropriate amount of one or more of the oligonucleotides is provided in one or more containers, or held on a substrate. An oligonucleotide primer or probe can be provided in an aqueous solution or as a freeze-dried or lyophilized powder, for instance. The container(s) in which the oligonucleotide(s) are supplied can be any conventional container that is capable of holding the supplied form, for instance, microfuge tubes, ampoules, or bottles. In some applications, pairs of primers are provided in pre-measured single use amounts in individual (typically disposable) tubes or equivalent containers. With such an arrangement, the sample to be tested for the presence of a SARS-CoV nucleic acid can be added to the individual tubes and amplification carried out directly.

The amount of each oligonucleotide supplied in the kit can be any appropriate amount, and can depend on the market to which the product is directed. For instance, if the kit is adapted for research or clinical use, the amount of each oligonucleotide primer provided would likely be an amount sufficient to prime several PCR amplification reactions. General guidelines for determining appropriate amounts can be found, for example, in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 2001; Ausubel et al. (eds.), *Short Protocols in Molecular Biology*, John Wiley and Sons, New York, N.Y., 1999; and Innis et al., *PCR Applications, Protocols for Functional Genomics*, Academic Press, Inc., San Diego, Calif., 1999. A kit can include more than two primers, in order to facilitate the amplification of a larger number of SARS-CoV nucleotide sequences.

In some embodiments, kits also include one or more reagents necessary to carry out in vitro amplification reactions, including DNA sample preparation reagents, appropriate buffers (for example, polymerase buffer), salts (for example, magnesium chloride), and deoxyribonucleotides (dNTP5).

Kits can include either labeled or unlabeled oligonucleotide primers and/or probes for use in detection of SARS-CoV nucleotide sequences. The appropriate sequences for such a probe will be any sequence that falls between the annealing sites of the two provided oligonucleotide primers,

such that the sequence that the probe is complementary to is amplified during the amplification reaction.

One or more control sequences for use in the amplification reactions also can be supplied in the kit. In other particular embodiments, the kit includes equipment, reagents, and instructions for extracting and/or purifying nucleotides from a sample.

Kits for the detection of SARS-CoV antigen include for instance at least one SARS-CoV antigen-specific binding agent (for example, a polyclonal or monoclonal antibody or antibody fragment). The kits may also include means for detecting antigen-specific binding agent complexes, for instance the specific binding agent may be detectably labeled. If the specific binding agent is not labeled, it may be detected by second antibodies or protein A, for example, which may also be provided in some kits in one or more separate containers. Such techniques are well known.

Another example of an assay kit provided herein is a recombinant SARS-CoV-specific polypeptide (or fragment thereof) as an antigen and an enzyme-conjugated anti-human antibody as a second antibody. Examples of such kits also can include one or more enzymatic substrates. Such kits can be used to test if a sample from a subject contains antibodies against a SARS-CoV-specific protein.

The subject matter of the present disclosure is further illustrated by the following non-limiting Examples.

EXAMPLES

Example 1

Isolation and Characterization of SARS-CoV

Virus Isolation and Ultrastructural Characterization

This example describes the original isolation and characterization of a new human coronavirus from patients with SARS.

A variety of clinical specimens (blood, serum, material from oropharyngeal swabs or washings, material from nasopharyngeal swabs, and tissues of major organs collected at autopsy) from patients meeting the case definition of SARS were sent to the Centers for Disease Control and Prevention (CDC) as part of the etiologic investigation of SARS. These samples were inoculated onto a number of continuous cell lines, including Vero E6, NCI-H292, MDCK, LLC-MK2, and B95-8 cells, and into suckling ICR mice by the intracranial and intraperitoneal routes. All cultures were observed daily for CPE. Maintenance medium was replenished at day seven, and cultures were terminated fourteen days after inoculation. Any cultures exhibiting identifiable CPE were subjected to several procedures to identify the cause of the effect. Suckling mice were observed daily for fourteen days, and any sick or dead mice were further tested by preparing a brain suspension that was filtered and subcultured. Mice that remained well after fourteen days were killed, and their test results were recorded as negative.

Two cell lines, Vero E6 cells and NCI-H292 cells, inoculated with oropharyngeal specimens from Patient 16 (a 46 year old male physician with an epidemiologic link to a hospital with multiple SARS patients) initially showed CPE (Table 1)

TABLE 1

Specimens from patients with SARS that were positive for SARS-CoV by one or more methods*.								
Patient No	Exposure and Setting	Age/Sex	Findings on Chest Radiograph	Hospitalization	Serologic Results	Specimen	Isolation	RT-PCR
1	Singapore, hospital	53 yr/F	Pneumonia	Yes	+	Nasal, oropharyngeal swabs	-	Not done
2†	Hong Kong, hotel	36 yr/F	Pneumonia	Yes	+	Nasal, swab	-	Not done
3	Hong Kong, hotel	22 yr/M	Pneumonia	Yes	+	Swab	-	-
4†	Hong Kong, hotel	39 yr/M	Pneumonia	Yes	+	Nasal, pharyngeal swab	-	-
5	Hong Kong, hotel	49 Yr/M	Pneumonia	Yes	Not done	Sputum	+	+
6‡	Hong Kong, hotel	46 yr/M	Pneumonia	Yes	+	Kidney, lung, broncho-alveolar lavage	+§	+
7	Vietnam, hospital	Adult/unknown	Pneumonia	Yes	-	Oropharyngeal wash	+	+
8	Vietnam, hospital	Adult/unknown	Pneumonia	Yes	-	Oropharyngeal wash	-	+
9	Vietnam, hospital	Adult/unknown	Pneumonia	Yes	-	Oropharyngeal wash	-	+
10	Vietnam, hospital	Adult/unknown	Pneumonia	Yes	-	Oropharyngeal wash	-	+
11	Vietnam, hospital	Adult/unknown	Pneumonia	Yes	-	Oropharyngeal wash	-	+
12	Vietnam, hospital	Adult/unknown	Pneumonia	Yes	-	Oropharyngeal wash	-	+
13	Vietnam, hospital	Adult/unknown	Pneumonia	Yes	-	Oropharyngeal wash	+	+
14	Vietnam, hospital	Adult/unknown	Pneumonia	Yes	-	Oropharyngeal wash	-	+
15	Vietnam, hospital	Adult/unknown	Pneumonia	Yes	-	Oropharyngeal wash	-	+
16	Vietnam, hospital	46 yr/M	Pneumonia	Yes	+	Nasal, oropharyngeal swabs	+¶	+
17	Canada, family	43 yr/M	Pneumonia	Yes	Not done	Lung, bone marrow	-	-
18	Taiwan, family	51 yr/F	Pneumonia	Yes	-	Sputum	-	+
19	Hong Kong, hotel	Adult/F	Pneumonia	Yes	+	Oropharyngeal wash	-	+

*Plus signs denote positive results, and minus signs negative results. The serologic and RT-PCR assays were not necessarily performed on samples obtained at the same time.

†This was a late specimen, antibody positive at first sample.

‡Travel included China, Hong Kong (hotel), and Hanoi (the patient was the index patient in the French Hospital).

§Isolation was from the kidney only.

¶Isolation was from the oropharyngeal only.

The CPE in the Vero E6 cells was first noted on the fifth day post-inoculation; it was focal, with cell rounding and a refractive appearance in the affected cells that was soon followed by cell detachment (FIG. 1A). The CPE spread quickly to involve the entire cell monolayer within 24 to 48 hours. Subculture of material after preparation of a master seed stock (used for subsequent antigen production) resulted in the rapid appearance of CPE, as noted above, and in complete destruction of the monolayer in the inoculated flasks within 48 hours. Similar CPE was also noted in four additional cultures: three cultures of respiratory specimens (two oropharyngeal washes and one sputum specimen) and one culture of a suspension of kidney tissue obtained at autopsy. In these specimens, the initial CPE was observed between day two and day four and, as noted above, the CPE rapidly progressed to involve the entire cell monolayer.

Tissue culture samples showing CPE were prepared for electron-microscopical examination. Negative-stain electron-microscopical specimens were prepared by drying culture supernatant, mixed 1:1 with 2.5% paraformaldehyde, onto Formvarcarbon-coated grids and staining with 2% methylamine tungstate. Thin-section electron-microscopical

specimens were prepared by fixing a washed cell pellet with 2.5% glutaraldehyde and embedding the cell pellet in epoxy resin. In addition, a master seed stock was prepared from the remaining culture supernatant and cells by freeze-thawing the culture flask, clarifying the thawed contents by centrifugation at 1000×g, and dispensing the supernatant into aliquots stored in gas phase over liquid nitrogen. The master seed stock was subcultured into 850-cm² roller bottles of Vero E6 cells for the preparation of formalin-fixed positive control cells for immunohistochemical analysis, mixed with normal Vero E6 cells, and gamma-irradiated for preparation of spot slides for IFA tests or extracted with detergent and gamma-irradiated for use as an ELISA antigen for antibody tests.

Examination of CPE-positive Vero E6 cells by thin-section electron microscopy revealed characteristic coronavirus particles within the cisternae of the rough endoplasmic reticulum and in vesicles (FIG. 2A) (Becker et al., *J. Virol.* 1:1019-27, 1967; Oshiro et al. *J. Gen. Virol.* 12:161-8, 1971). Extracellular particles were found in large clusters and adhering to the surface of the plasma membrane. Negative-stain electron microscopy identified coronavirus particles, 80 to 140 nm in diameter, with 20- to 40-nm complex surface projections

surrounding the periphery (FIG. 2B). Hemagglutinin esterase-type glycoprotein projections were not seen.

The isolation and growth of a human-derived coronavirus in Vero E6 cells were unexpected. The previously known human coronaviruses are notably fastidious, preferring select cell lines, organ culture, or suckling mice for propagation. The only human or animal coronavirus which has been shown to grow in Vero E6 cells is PEDV, and it requires the addition of trypsin to culture medium for growth in the cells. Moreover, PEDV adapted to growth in Vero E6 cells results in a strikingly different CPE, with cytoplasmic vacuoles and the formation of large syncytia. Syncytial cells were only observed occasionally in monolayers of Vero E6 cells infected with the SARS-CoV; they clearly do not represent the dominant CPE.

Reverse Transcription-Polymerase Chain Reaction and Sequencing

For RT-PCR assays, cell-culture supernatants were placed in lysis buffer. RNA extracts were prepared from 100 μ l of each specimen (or culture supernatant) with the automated NucliSens extraction system (bioMérieux, Durham, N.C.). Initially, degenerate, inosine-containing primers IN-2 (+) 5'GGGTTGGGACTA TCCTAAGTGTGA3' (SEQ ID NO: 34) and IN-4(-) 5'TAACACACAACICCATCA TCA3' (SEQ ID NO: 35) were designed to anneal to sites encoding conserved amino acid motifs that were identified on the basis of alignments of available coronavirus ORF1a, ORF1b, S, HE, M, and N gene sequences. Additional, SARS-specific, primers Cor-p-F2 (+) 5'CTAACATGCTTAGGATAATGG3' (SEQ ID NO: 13), Cor-p-F3 (+) 5'GCCTCTCTTGTCT-TGCTCGC3' (SEQ ID NO: 14), and Cor-p-R1 (-) 5' CAG-GTAAAGCGTAAACTCATC3' (SEQ ID NO: 15) were designed as sequences were generated from RT-PCR products amplified with the degenerate primers. These SARS-specific primers were used to test patient specimens for SARS (see below). Primers used for specific amplification of human metapneumovirus have been described by Falsey et al. (*J. Infect. Dis.* 87:785-90, 2003).

For RT-PCR products of less than 3 kb, cDNA was synthesized in a 20 μ l reaction mixture containing 500 ng of RNA, 200 U of Superscript™ II reverse transcriptase (Invitrogen Life Technologies, Carlsbad, Calif.), 40 U of RNasin (Promega Corp., Madison, Wis.), 100 mM each dNTP (Roche Molecular Biochemicals, Indianapolis, Ind.), 4 μ l of 5 \times reaction buffer (Invitrogen Life Technologies, Carlsbad, Calif.), and 200 pmol of the reverse primer. The reaction mixture, except for the reverse transcriptase, was heated to 70° C. for 2 minutes, cooled to 4° C. for 5 minutes and then heated to 42° C. in a thermocycler. The mixture was held at 42° C. for 4 minutes, and then the reverse transcriptase was added, and the reactions were incubated at 42° C. for 45 minutes. Two microliters of the cDNA reaction was used in a 50 μ l PCR reaction containing 67 mM Tris-HCl (pH 8.8), 1 mM each primer, 17 mM ammonium sulfate, 6 mM EDTA, 2 mM MgCl₂, 200 mM each dNTP, and 2.5 U of Taq DNA polymerase (Roche Molecular Biochemicals, Indianapolis, Ind.). The thermocycler program for the PCR consisted of 40 cycles of denaturation at 95° C. for 30 seconds, annealing at 42° C. for 30 seconds, and extension at 65° C. for 30 seconds. For SARS-CoV-specific primers, the annealing temperature was increased to 55° C.

For amplification of fragments longer than 3 kb, regions of the genome between sections of known sequence were amplified by means of a long RT-PCR protocol and SARS-CoV-specific primers. First-strand cDNA synthesis was performed at 42° C. or 50° C. using Superscript™ II RNase H reverse

transcriptase (Invitrogen Life Technologies, Carlsbad, Calif.) according to the manufacturer's instructions with minor modifications. Coronavirus-specific primers (500 ng) and SARS-CoV RNA (350 ng) were combined with the PCR Nucleotide Mix (Roche Molecular Biochemicals, Indianapolis, Ind.), heated for 1 minute at 94° C., and cooled to 4° C. in a thermocycler. The 5 \times first-strand buffer, dithiothreitol (Invitrogen Life Technologies, Carlsbad, Calif.), and Protector RNase Inhibitor (Roche Molecular Biochemicals, Indianapolis, Ind.) were added, and the samples were incubated at 42° C. or 50° C. for 2 minutes. After reverse transcriptase (200 U) was added, the samples were incubated at 42° C. or 50° C. for 1.5 to 2 hours. Samples were inactivated at 70° C. for 15 minutes and subsequently treated with 2 U of RNase H (Roche Molecular Biochemicals, Indianapolis, Ind.) at 37° C. for 30 minutes. Long RT-PCR amplification of 5- to 8-kb fragments was performed using Taq Plus Precision (Stratagene, La Jolla, Calif.) and AmpliWax PCR Gem 100 beads (Applied Biosystems; Foster City, Calif.) for "hot start" PCR with the following thermocycling parameters: denaturation at 94° C. for 1 minute followed by 35 cycles of 94° C. for 30 seconds, 55° C. for 30 seconds, an increase of 0.4 degrees per second up to 72° C., and 72° C. for 7 to 10 minutes, with a final extension at 72° C. for 10 minutes. RT-PCR products were separated by electrophoresis on 0.9% agarose TAE gels and purified by use of a QIAquick Gel Extraction Kit (Qiagen, Inc., Santa Clarita, Calif.).

In all cases, the RT-PCR products were gel-isolated and purified for sequencing by means of a QIAquick Gel Extraction kit (Qiagen, Inc., Santa Clarita, Calif.). Both strands were sequenced by automated methods, using fluorescent dideoxy-chain terminators (Applied Biosystems; Foster City, Calif.).

The sequence of the leader was obtained from the subgenomic mRNA coding for the N gene and from the 5' terminus of genomic RNA. The 5' rapid amplification of cDNA ends (RACE) technique (Harcourt et al., *Virology* 271:334-49, 2000) was used with reverse primers specific for the N gene or for the 5' untranslated region. RACE products were either sequenced directly or were cloned into a plasmid vector before sequencing. A primer that was specific for the leader of SARS-CoV was used to amplify the region between the 5'-terminus of the genome and known sequences in the rep gene. The 3'-terminus of the genome was amplified for sequencing by use of an oligo-(dT) primer and primers specific for the N gene.

Once the complete SARS-CoV genomic sequence had been determined, it was confirmed by sequencing a series of independently amplified RT-PCR products spanning the entire genome. Positive- and negative-sense sequencing primers, at intervals of approximately 300 nt, were used to generate a confirmatory sequence with an average redundancy of 9.1. The confirmatory sequence was identical to the original sequence. The genomic sequence (SEQ ID NO: 1) was published in the GenBank sequence database (Accession No. AY278741) on Apr. 21, 2003.

Sequence Analysis

Predicted amino acid sequences were compared with those from reference viruses representing each species for which complete genomic sequence information was available: group 1 representatives included human coronavirus 229E (GenBank Accession No. AF304460), porcine epidemic diarrhea virus (GenBank Accession No. AF353511), and transmissible gastroenteritis virus (GenBank Accession No. AF271965); group 2 representatives included bovine coronavirus (GenBank Accession No. AF220295) and mouse hepatitis virus (GenBank Accession No. AF201929); group 3

was represented by infectious bronchitis virus (GenBank Accession No. M95169). Sequences for representative strains of other coronavirus species for which partial sequence information was available were included for some of the structural protein comparisons: group 1 representative strains included canine coronavirus (GenBank Accession No. D13096), feline coronavirus (GenBank Accession No. AY204704), and porcine respiratory coronavirus (GenBank Accession No. Z24675); and group 2 representatives included three strains of human coronavirus OC43 (GenBank Accession Nos. M76373, L14643 and M93390), porcine hemagglutinating encephalomyelitis virus (GenBank Accession No. AY078417), and rat coronavirus (GenBank Accession No. AF207551).

Partial nucleotide sequences of the polymerase gene were aligned with published coronavirus sequences, using CLUSTAL W for Unix (version 1.7; Thompson et al., *Nucleic Acids Res.* 22:4673-80, 1994). Phylogenetic trees were computed by maximum parsimony, distance, and maximum likelihood-based criteria analysis with PAUP (version 4.0.d10; Swofford ed., *Phylogenetic Analysis using Parsimony and other Methods*, Sinauer Associates, Sunderland, Mass.). When compared with other human and animal coronaviruses, the nucleotide and deduced amino acid sequence from this region had similarity scores ranging from 0.56 to 0.63 and from 0.57 to 0.74, respectively. The highest sequence simi-

Gonnet protein comparison matrix. The resulting trees were adjusted for final output by using treeool version 2.0.1. Uncorrected pairwise distances were calculated from the unaligned sequences by using the Distances program from the Wisconsin Sequence Analysis Package, version 10.2 (Accelrys, Burlington, Mass.). Distances were converted to percent identity by subtracting from 100. The amino acid sequences for three well-defined enzymatic proteins encoded by the rep gene and the four major structural proteins of SARS-CoV were compared with those from representative viruses for each of the species of coronavirus for which complete genomic sequence information was available (FIG. 4, Table 2). The topologies of the resulting phylograms are remarkably similar (FIG. 4). For each protein analyzed, the species formed monophyletic clusters consistent with the established taxonomic groups. In all cases, SARS-CoV sequences segregated into a fourth, well-resolved branch. These clusters were supported by bootstrap values above 90% (1000 replicates). Consistent with pairwise comparisons between the previously characterized coronavirus species (Table 2), there was greater sequence conservation in the enzymatic proteins (3CL^{PRO}, polymerase (POL), and helicase (HEL)) than among the structural proteins (S, E, M, and N). These results indicate that SARS-CoV is not closely related to any of the previously characterized coronaviruses and forms a distinct group within the genus *Coronavirus*.

TABLE 2

Pairwise amino acid identities of coronavirus proteins.								
Group	Virus	3CLPRO	POL	HEL	S	E	M	N
Pairwise Amino Acid Identity (Percent)								
G1	HCoV-229E	40.1	58.8	59.7	23.9	22.7	28.8	23.0
	PEDV	44.4	59.5	61.7	21.7	17.6	31.8	22.6
	TGEV	44.0	59.4	61.2	20.6	22.4	30.0	25.6
G2	BCoV	48.8	66.3	68.3	27.1	20.0	39.7	31.9
	MHV	49.2	66.5	67.3	26.5	21.1	39.0	33.0
G3	IBV	41.3	62.5	58.6	21.8	18.4	27.2	24.0
Predicted Protein Length (aa)								
	SARS-CoV	306	932	601	1255	76	221	422
	CoV Range	302-307	923-940	506-600	1173-1452	76-108	225-262	377-454

ilarity was obtained with group II coronaviruses. The maximum-parsimony tree obtained from the nucleotide-sequence alignment is shown in FIG. 3. Bootstrap analyses of the internal nodes at the internal branches of the tree provided strong evidence that the SARS-CoV is genetically distinct from other known coronaviruses.

Microarray analyses (using a long oligonucleotide DNA microarray with array elements derived from highly conserved regions within viral families) of samples from infected and uninfected cell cultures gave a positive signal for a group of eight oligonucleotides derived from two virus families: Coronaviridae and Astroviridae (Wang et al., PNAS 99:15687-92, 2002). All of the astroviruses and two of the coronavirus oligonucleotides share a consensus sequence motif that maps to the extreme 3'-end of astroviruses and two members of the coronavirus family: avian infectious bronchitis and turkey coronavirus (Jonassen et al., *J. Gen. Virol.* 79:715-8, 1998). Results were consistent with the identity of the isolate as a coronavirus.

Additional sequence alignments and neighbor-joining trees were generated by using ClustalX (Thompson et al., *Nucleic Acids Res.* 25:4876-82, 1997), version 1.83, with the

Example 2

Detection of SARS-CoV in a Subject

This example demonstrates the detection of SARS-CoV in patient specimens using SARS-CoV-specific primers.

The SARS-specific primers Cor-p-F2 (SEQ ID NO: 13), Cor-p-F3 (SEQ ID NO: 14) and Cor-p-R1 (SEQ ID NO: 15) were used to test patient specimens for SARS. One primer for each set was 5'-end-labeled with 6-FAM to facilitate GeneScan analysis. One-step amplification reactions were performed with the Access RT-PCR System (Promega, Madison, Wis.) as described by Falsey et al., *J. Infect. Dis.* 87:785-90, 2003. Positive and negative RT-PCR controls, containing standardized viral RNA extracts, and nuclease-free water were included in each run. Amplified 6-FAM-labeled products were analyzed by capillary electrophoresis on an ABI 3100 Prism Genetic Analyzer with GeneScan software (version 3.1.2; Applied Biosystems; Foster City, Calif.). Specimens were considered positive for SARS-CoV if the amplification products were within one nucleotide of the expected product size (368 nucleotides for Cor-p-F2 or Cor-p-R1 and

348 nucleotides for Cor-p-F3 or Cor-p-R1) for both specific primer sets, as confirmed by a second PCR reaction from another aliquot of RNA extract in a separate laboratory. Where DNA yield was sufficient, the amplified products were also sequenced. Additionally, as described above, microarray-based detection of SARS-CoV in patient specimens was carried out (Wang et al., *PNAS* 99:15687-92, 2002 and Bohlander et al., *Genomics* 13:1322-24, 1992).

Example 3

Immunohistochemical and Histopathological Analysis, and Electron-Microscopical Analysis of Bronchoalveolar Lavage Fluid

This example illustrates immunohistochemical, histopathological and electron-microscopical analysis of Vero E6 cells infected with the SARS-CoV and tissue samples from SARS patients.

Formalin-fixed, paraffin-embedded Vero E6 cells infected with the SARS-CoV and tissues obtained from patients with SARS were stained with hematoxylin and eosin and various immunohistochemical stains. Immunohistochemical assays were based on a method described previously for hantavirus (Zaki et al., *Amer. J. Pathol.* 146:552-79, 1995). Briefly, 4- μ m sections were deparaffinized, rehydrated, and digested in Proteinase K for 15 minutes. Slides were then incubated for 60 minutes at room temperature with monoclonal antibodies, polyclonal antiserum or ascitic fluids derived from animal species with reactivities to various known coronaviruses, and with a convalescent-phase serum specimen from a patient with SARS.

Optimal dilutions of the primary antibodies were determined by titration experiments with coronavirus-infected cells from patients with SARS and with noninfected cells or, when available, with concentrations recommended by the manufacturers. After sequential application of the appropriate biotinylated link antibody, avidin-alkaline phosphatase complex, and naphthol-fast red substrate, sections were counterstained in Mayer's hematoxylin and mounted with aqueous mounting medium. The following antibody and tissue controls were used: serum specimens from noninfected animals, various coronavirus-infected cell cultures and animal tissues, noninfected cell cultures, and normal human and animal tissues. Tissues from patients were also tested by immunohistochemical assays for various other viral and bacterial pulmonary pathogens. In addition, a BAL specimen was available from one patient for thin-section electron-microscopical evaluation.

Lung tissues were obtained from the autopsy of three patients and by open lung biopsy of one patient, 14-19 days following onset of SARS symptoms. Confirmatory laboratory evidence of infection with coronavirus was available for two patients (patients 6 and 17) and included PCR amplification of coronavirus nucleic acids from tissues, viral isolation from BAL fluid or detection of serum antibodies reactive with coronavirus (Table 1). For two patients, no samples were available for molecular, cell culture, or serological analysis; however, both patients met the CDC definition for probable SARS cases and had strong epidemiologic links with laboratory-confirmed SARS cases. Histopathologic evaluation of lung tissues of the four patients showed diffuse alveolar damage at various levels of progression and severity. Changes included hyaline membrane formation, interstitial mononuclear inflammatory infiltrates, and desquamation of pneumocytes in alveolar spaces (FIG. 5A). Other findings identified in some patients included focal intraalveolar

hemorrhage, necrotic inflammatory debris in small airways, and organizing pneumonia. Multinucleated syncytial cells were identified in the intraalveolar spaces of two patients who died 14 and 17 days, respectively, after onset of illness. These cells contained abundant vacuolated cytoplasm with cleaved and convoluted nuclei. No obvious intranuclear or intracytoplasmic viral inclusions were identified (FIG. 5B), and electron-microscopical examination of a limited number of these syncytial cells revealed no coronavirus particles. No definitive immunostaining was identified in tissues from SARS patients with the use of a battery of immunohistochemical stains reactive with coronaviruses from antigenic groups I, II, and III. In addition, no staining of patient tissues was identified with the use of immunohistochemical stains for influenza viruses A and B, adenoviruses, Hendra and Nipah viruses, human metapneumovirus, respiratory syncytial virus, measles virus, *Mycoplasma pneumoniae*, and *Chlamydia pneumoniae*.

Evaluation of Vero E6 cells infected with coronavirus isolated from a patient with SARS revealed viral CPE that included occasional multinucleated syncytial cells but no obvious viral inclusions (FIG. 5C). Immunohistochemical assays with various antibodies reactive with coronaviruses from antigenic group I, including HCoV-229E, FIPV and TGEV, and with an immune serum specimen from a patient with SARS, demonstrated strong cytoplasmic and membranous staining of infected cells (FIG. 5C and Table 3); however, cross-reactivity with the same immune human serum sample and FIPV antigen was not observed. No staining was identified with any of several monoclonal or polyclonal antibodies reactive with coronaviruses in antigenic group II (HCoV-OC43, BCoV and MHV) or group III (TCoV and IBV-Avian). Electron microscopical examination of a BAL fluid from one patient revealed many coronavirus-infected cells (FIGS. 6A-B).

TABLE 3

Immunohistochemical reactivities of various polyclonal group I anti-coronavirus reference antiserum samples with a coronavirus isolated from a patient with SARS and with selected antigenic group I coronaviruses.

Antiserum	Immunohistochemical reactivity of antiserum with coronavirus-infected culture cells		
	SARS-CoV (Vero E6)	HCoV-229E (mouse 3T3-hAPN)	FIPV-1 (BHK-fAPN)
Convalescent-phase SARS (patient 3)	+	+	-
Guinea pig anti-HCoV-229E	+	+	-
Rabbit anti-HCoV-229E	+	+	+
Feline anti-FIPV-1	+	+	+
Porcine anti-TGEV	+	-	+

Example 4

SARS-CoV Serologic Analysis

This example illustrates representative methods of performing serological analysis of SARS-CoV.

Spot slides were prepared by applying 15 μ l of the suspension of gamma-irradiated mixed infected and noninfected cells onto 12-well Teflon-coated slides. Slides were allowed

to air dry before being fixed in acetone. Slides were then stored at -70° C. until used for IFA tests (Wulff and Lange, *Bull. WHO* 52:429-36, 1975). An ELISA antigen was prepared by detergent extraction and subsequent gamma irradiation of infected Vero E6 cells (Ksiazek et al., *J. Infect. Dis.* 179 (suppl. 1):S191-8, 1999). The optimal dilution (1:1000) for the use of this antigen was determined by checkerboard titration against SARS patient serum from the convalescent phase; a control antigen, similarly prepared from uninfected Vero E6 cells, was used to control for specific reactivity of tested sera. The conjugates used were goat antihuman IgG, IgA, and IgM conjugated to fluorescein isothiocyanate and horseradish peroxidase (Kirkegaard and Perry, Gaithersburg, Md.), for the IFA test and ELISA, respectively. Specificity and cross-reactivity of a variety of serum samples to the newly identified virus were evaluated by using the tests described herein. For this evaluation, serum from SARS patients in Singapore, Bangkok and Hong Kong was used, along with serum from healthy blood donors from the CDC serum bank and from persons infected with known human coronavirus (human coronaviruses OC43 and 229E) (samples provided by E. Walsh and A. Falsey, University of Rochester School of Medicine and Dentistry, Rochester, N.Y.).

Spot slides with infected cells reacted with serum from patients with probable SARS in the convalescent phase (FIG. 1B). Screening of a panel of serum from patients with suspected SARS from Hong Kong, Bangkok, Singapore as well as the United States showed a high level of specific reaction with infected cells, and conversion from negative to positive reactivity or diagnostic rises in the IFA test by a factor of four. Similarly, tests of these same serum samples with the ELISA antigen showed high specific signal in the convalescent-phase samples and conversion from negative to positive antibody reactivity or diagnostic increases in titer (Table 4).

TABLE 4

Results of serological testing with both IFA assay and ELISA in SARS patients tested against the newly isolated human coronavirus.				
Source	Serum No.	Days After Onset	ELISA Titer*	IFA Titer*
Hong Kong	1.1	4	<100	<25
Hong Kong	1.2	13	\geq 6400	1600
Hong Kong	2.1	11	400	100
Hong Kong	2.2	16	1600	200
Hong Kong	3.1	7	<100	<25
Hong Kong	3.2	17	\geq 6400	800
Hong Kong	4.1	8	<100	<25
Hong Kong	4.2	13	1600	50
Hong Kong	5.1	10	100	<25
Hong Kong	5.2	17	\geq 6400	1600
Hong Kong	6.1	12	1600	200
Hong Kong	6.2	20	\geq 6400	6400
Hong Kong	7.1	17	400	50
Hong Kong	7.2	24	\geq 6400	3200
Hong Kong	8.1	3	<100	<25
Hong Kong	8.2	15	\geq 6400	200
Hong Kong (Hanoi)	9.1	5	<100	<25
Hong Kong	9.2	11	\geq 6400	1600
Bangkok	1.1	2	<100	<25
Bangkok	1.2	4	<100	<25
Bangkok	1.3	7	<100	<25
Bangkok	1.4	15	1600	200
United States	1.1	2	<100	<25
United States	1.2	6	400	50
United States	1.3	13	\geq 6400	800
Singapore	1.1	2	100	<25
Singapore	1.2	11	\geq 6400	800
Singapore	2.1	6	100	<25
Singapore	2.2	25	\geq 6400	400

TABLE 4-continued

Results of serological testing with both IFA assay and ELISA in SARS patients tested against the newly isolated human coronavirus.				
Source	Serum No.	Days After Onset	ELISA Titer*	IFA Titer*
Singapore	3.1	6	100	<25
Singapore	3.2	14	\geq 6400	400
Singapore	4.1	5	100	<25
Singapore	4.2	16	1600	400

*Reciprocal of dilution

Information from the limited numbers of samples tested suggests that antibody is first detectable by IFA assay and ELISA between one and two weeks after the onset of symptoms in the patient. IFA testing and ELISA of a panel of 384 randomly selected serum samples (from U.S. blood donors) were negative for antibodies to the new coronavirus, with the exception of one specimen that had minimal reactivity on ELISA. A panel of paired human serum samples with diagnostic increases (by a factor of four or more) in antibody (with very high titers to the homologous viral antigen in the convalescent-phase serum) to the two known human coronaviruses, OC43 (13 pairs) and 229E (14 pairs), showed no reactivity in either acute- or convalescent-phase serum with the newly isolated coronavirus by either the IFA test or the ELISA.

Example 5

Poly(A)⁺RNA Isolation and Northern Hybridization

This example illustrates a representative method of Northern hybridization to detect SARS-CoV messages in Vero E6 cells.

Total RNA from infected or uninfected Vero E6 cells was isolated with Trizol reagent (Invitrogen Life Technologies, Carlsbad, Calif.) used according to the manufacturer's recommendations. Poly(A)⁺RNA was isolated from total RNA by use of the Oligotex Direct mRNA Kit (Qiagen, Inc., Santa Clarita, Calif.), following the instructions for the batch protocol, followed by ethanol precipitation. RNA isolated from 1 cm² of cells was separated by electrophoresis on a 0.9% agarose gel containing 3.7% formaldehyde, followed by partial alkaline hydrolysis (Ausubel et al. eds. *Current Protocols in Molecular Biology*, vol. 1, John Wiley & Sons, Inc., NY, N.Y., Ch. 4.9, 1996). RNA was transferred to a nylon membrane (Roche Molecular Biochemicals, Indianapolis, Ind.) by vacuum blotting (Bio-Rad, Hercules, Calif.) and fixed by UV cross-linking. The DNA template for probe synthesis was generated by RT-PCR amplification of SARS-CoV nt 29,083 to 29,608 (SEQ ID NO: 1), by using a reverse primer containing a T7 RNA polymerase promoter to facilitate generation of a negative-sense riboprobe. In vitro transcription of the digoxigenin-labeled riboprobe, hybridization, and detection of the bands were carried out with the digoxigenin system by using manufacturer's recommended procedures (Roche Molecular Biochemicals, Indianapolis, Ind.). Signals were visualized by chemiluminescence and detected with x-ray film.

Example 6

SARS-CoV Genome Organization

This example illustrates the genomic organization of the SARS-CoV genome, including the location of SARS-CoV ORFs.

The genome of SARS-CoV is a 29,727-nucleotide, polyadenylated RNA, and 41% of the residues are G or C (range for published coronavirus complete genome sequences, 37% to 42%). The genomic organization is typical of coronaviruses, having the characteristic gene order [5'-replicase (rep), spike (S), envelope (E), membrane (M), nucleocapsid (N)-3'] and short untranslated regions at both termini (FIG. 7A, Table 5). The SARS-CoV rep gene, which comprises approximately two-thirds of the genome, encodes two polyproteins (encoded by ORF1a and ORF1b) that undergo co-translational proteolytic processing. There are four ORFs downstream of rep that encode the structural proteins, S, E, M, and N, which are common to all known coronaviruses. The hemagglutinin-esterase gene, which is present between ORF1b and S in group 2 and some group 3 coronaviruses (Lai and Holmes, in *Fields Virology*, eds. Knipe and Howley, Lippincott Williams and Wilkins, New York, 4th edition, 2001, Ch. 35), was not found in SARS-CoV.

Coronaviruses also encode a number of non-structural proteins that are located between S and E, between M and N, or downstream of N. These non-structural proteins, which vary widely among the different coronavirus species, are of unknown function and are dispensable for virus replication (Lai and Holmes, in *Fields Virology*, eds. Knipe and Howley, Lippincott Williams and Wilkins, New York, 4th edition, 2001, Ch. 35). The genome of SARS-CoV contains ORFs for five non-structural proteins of greater than 50 amino acids (FIG. 7B, Table 5). Two overlapping ORFs encoding proteins of 274 and 154 amino acids (termed X1 (SEQ ID NO: 5) and X2 (SEQ ID NO: 6), respectively) are located between S (SEQ ID NO: 4) and E (SEQ ID NO: 7). Three additional non-structural genes, X3 (SEQ ID NO: 9), X4 (SEQ ID NO: 10), and X5 (SEQ ID NO: 11) (encoding proteins of 63, 122, and 84 amino acids, respectively), are located between M (SEQ ID NO: 8) and N (SEQ ID NO: 12). In addition to the five ORFs encoding the non-structural proteins described above, there are also two smaller ORFs between M and N, encoding proteins of less than 50 amino acids. Searches of the GenBank database (BLAST and FastA) indicated that there is no significant sequence similarity between these non-structural proteins of SARS-CoV and any other proteins.

The coronavirus rep gene products are translated from genomic RNA, but the remaining viral proteins are translated from subgenomic mRNAs that form a 3'-coterminal nested set, each with a 5'-end derived from the genomic 5'-leader sequence. The coronavirus subgenomic mRNAs are synthesized through a discontinuous transcription process, the mechanism of which has not been unequivocally established (Lai and Holmes, in *Fields Virology*, eds. Knipe and Howley, Lippincott Williams and Wilkins, New York, 4th edition, 2001, Ch. 35; Sawicki and Sawicki, *Adv. Exp. Med. Biol.* 440:215-19, 1998). The SARS-CoV leader sequence was mapped by comparing the sequence of 5'-RACE products synthesized from the N gene mRNA with those synthesized from genomic RNA. A sequence, AAACGAAC (nucleotides 65-72 of SEQ ID NO: 1), was identified immediately upstream of the site where the N gene mRNA and genomic sequences diverged. This sequence was also present upstream of ORF1a and immediately upstream of five other ORFs (Table 5), suggesting that it functions as the conserved core of the transcriptional regulatory sequence (TRS).

In addition to the site at the 5'-terminus of the genome, the TRS conserved core sequence appears six times in the remainder of the genome. The positions of the TRS in the genome of SARS-CoV predict that subgenomic mRNAs of 8.3, 4.5, 3.4, 2.5, 2.0, and 1.7 kb, not including the poly(A) tail, should be produced (FIGS. 7A-B, Table 5). At least five subgenomic mRNAs were detected by Northern hybridization of RNA from SARS-CoV-infected cells, using a probe derived from the 3'-untranslated region (FIG. 7C). The calculated sizes of the five predominant bands correspond to the sizes of five of the predicted subgenomic mRNAs of SARS-CoV; the possibility that other, low-abundance mRNAs are present cannot be excluded. By analogy with other coronaviruses (Lai and Holmes, in *Fields Virology*, eds. Knipe and Howley, Lippincott Williams and Wilkins, New York, 4th edition, 2001, Ch. 35), the 8.3-kb and 1.7-kb subgenomic mRNAs are monocistronic, directing translation of S and N, respectively, whereas multiple proteins are translated from the 4.5-kb (X1, X2, and E), 3.4-kb (M and X3), and 2.5-kb (X4 and X5) mRNAs. A consensus TRS is not found directly upstream of the ORF encoding the predicted E protein, and a monocistronic mRNA that would be predicted to code for E could not be clearly identified by Northern blot analysis. It is possible that the 3.6-kb band contains more than one mRNA species or that the monocistronic mRNA for E is a low-abundance message.

TABLE 5

Locations of SARS-CoV ORFs and sizes of proteins and mRNAs						
Genome Location				Predicted Size		
ORF	TRS ^a	ORF Start	ORF End	Protein (aa)	mRNA (nt) ^b	
1a	72	265	13,398	4,378	29,727	
1b			13,398	2,695		
S	21,491	21,492	25,256	1,255	8,308 ^c	
X1	25,265	25,268	26,089	274	4,534 ^c	
X2		25,689	26,150	154		
E		26,117	26,344	76		
M	26,353	26,398	27,060	221	3,446 ^c	
X3		27,074	27,262	63		
X4	27,272	27,273	27,638	122	2,527 ^c	
X5	27,778	27,864	28,115	84	2,021 ^d	
N	28,111	28,120	29,385	422	1,688 ^c	

^aThe location is the 3'-most nucleotide in the consensus TRS, AAACGAAC.

^bNot including poly(A). Predicted size is based on the position of the conserved TRS.

^cCorresponding mRNA detected by Northern blot analysis (FIG. 7C)

^dNo mRNA corresponding to utilization of this consensus TRS was detected by Northern blot analysis (FIG. 7C)

Example 7

Real-Time RT-PCR Assay for SARS-CoV Detection

This example demonstrates the use of SARS-CoV-specific primers and probes in a real-time RT-PCR assay to detect SARS-CoV in patient specimens.

A variant of the real-time format, based on TaqMan probe hydrolysis technology (Applied Biosystems, Foster City, Calif.), was used to analyze a total of 340 clinical specimens collected from 246 persons with confirmed or suspected SARS-CoV infection. Specimens included oro- and nasopharyngeal swabs (dry and in viral transport media), sputa, nasal aspirates and washes, BAL, and lung tissue specimens collected at autopsy.

Nucleic Acid Extraction

SARS-CoV nucleic acids were recovered from clinical specimens using the automated NucliSens extraction system (bioMérieux, Durham, N.C.). Following manufacturer's instructions, specimens received in NucliSens lysis buffer were incubated at 37° C. for 30 min with intermittent mixing, and 50 µL of silica suspension, provided in the extraction kit, was added and mixed. The contents of the tube were then transferred to a nucleic acid extraction cartridge and processed on an extractor workstation. Approximately 40-50 µL of total nucleic acid eluate was recovered into nuclease-free vials and either tested immediately or stored at -70° C.

Primers and Probes

Multiple primer and probe sets were designed from the SARS-CoV polymerase 1b (nucleic acid 13,398 to 21,482 of SEQ ID NO: 1) and nucleocapsid gene (nucleic acid 28,120 to 29,385 of SEQ ID NO: 1) sequences by using Primer Express software version 1.5 or 2.0.0 (Applied Biosystems, Foster City, Calif.) with the following default settings: primer melting temperature (T_M) set at 60° C.; probe T_M set at 10° C. greater than the primers at approximately 70° C.; and no guanidine residues permitted at the 5' probe termini. All primers and probes were synthesized by standard phosphoramidite chemistry techniques. TaqMan probes were labeled at the 5'-end with the reporter 6-FAM and at the 3'-end with the quencher Blackhole Quencher 1 (Biosearch Technologies, Inc., Novato, Calif.). Optimal primer and probe concentrations were determined by cross-titration of serial twofold dilutions of each primer against a constant amount of purified SARS-CoV RNA. Primer and probe concentrations that gave the highest amplification efficiencies were selected for further study (Table 6).

0.25 µL each of 50 µM forward and reverse primers, 6.125 µL of nuclease-free water, and 5 µL of nucleic acid extract. Amplification was carried out in 96-well plates on an iCycler iQ Real-Time Detection System (Bio-Rad, Hercules, Calif.). Thermocycling conditions consisted of 30 minutes at 48° C. for reverse transcription, 10 minutes at 95° C. for activation of the AmpliTaq Gold DNA polymerase, and 45 cycles of 15 seconds at 95° C. and 1 minute at 60° C. Each run included one SARS-CoV genomic template control and at least two no-template controls for the extraction (to check for contamination during sample processing) and one no-template control for the PCR-amplification step. As a control for PCR inhibitors, and to monitor nucleic acid extraction efficiency, each sample was tested by real-time RT-PCR for the presence of the human ribonuclease (RNase) P gene (GenBank Accession No. NM_006413) by using the following primers and probe: forward primer 5'-AGATTGGACCTGCGAGCG-3' (SEQ ID NO: 36); reverse primer 5'-GAGCGGCTGTCTC-CACAAGT-3' (SEQ ID NO: 37); probe 5'-TTCTGACC TGAAGGCTCTGCGCG-3' (SEQ ID NO: 38). The assay reaction was performed identically to that described above except that primer concentrations used were 30 µM each. Fluorescence measurements were taken and the threshold cycle (C_T) value for each sample was calculated by determining the point at which fluorescence exceeded a threshold limit set at the mean plus 10 standard deviations above the baseline. A test result was considered positive if two or more of the SARS genomic targets showed positive results ($C_T \leq 45$ cycles) and all positive and negative control reactions gave expected values.

While this disclosure has been described with an emphasis upon preferred embodiments, it will be obvious to those of

TABLE 6

Primers and probes used for real-time RT-PCR assays ^a			
Assay ID	Primer/probe	Sequence	Genomic Region
Primary diagnostic assay			
SARS1	F	CAITGTGTGGCGCTCACTATAT (SEQ ID NO: 16)	RNA Pol
	R	GACACTATTAGCATAAGCAGTTGTAGCA (SEQ ID NO: 17)	
	P	TTAAACCAGGTGGAACATCATCCGGTG (SEQ ID NO: 18)	
SARS2	F	GGAGCCTTGAATACACCCAAAG (SEQ ID NO: 19)	Nucleocapsid
	R	GCACGGTGGCAGCATTG (SEQ ID NO: 20)	
	P	CCACATTGGCACCCGCAATCC (SEQ ID NO: 21)	
SARS3	F	CAAACATTGGCCGCAAAT (SEQ ID NO: 22)	Nucleocapsid
	R	CAATGCGTGACATTCCAAAGA (SEQ ID NO: 23)	
	P	CACAATTGCTCCAAGTGCCTCTGCA (SEQ ID NO: 24)	
To confirm positive results			
N3	F	GAAGTACCAITCTGGGGCTGAG (SEQ ID NO: 25)	Nucleocapsid
	R	CCGAAGAGCTACCCGACG (SEQ ID NO: 26)	
	P	CTCTTTCATTTTGGCCGTACCCACCAC (SEQ ID NO: 27)	
3'-NTR	F	AGCTCTCCCTAGCATTATTCAGT (SEQ ID NO: 28)	3'-NTR
	R	CACCACATTTTCATCGAGGC (SEQ ID NO: 29)	
	P	TACCCTCGATCGTACTCCGCGT (SEQ ID NO: 30)	
M	F	TGTAGGCACATGATTCAGGTTTG (SEQ ID NO: 31)	M protein
	R	CGGCGTGGTCTGTATTAAITTA (SEQ ID NO: 32)	
	P	CTGCATACAACCGCTACCGTATTGGAA (SEQ ID NO: 33)	

^aRT-PCR, reverse transcription-polymerase chain reaction; F, forward primer; R, reverse primer; P, probe; NTR, nontranslated region.

Real-Time RT-PCR Assay

The real-time RT-PCR assay was performed by using the Real-Time One-Step RT-PCR Master Mix (Applied Biosystems, Foster City, Calif.). Each 25-µL reaction mixture contained 12.5 µL of 2× Master Mix, 0.625 µL of the 40× MultiScribe and RNase Inhibitor mix, 0.25 µL of 10 µM probe,

ordinary skill in the art that variations and equivalents of the preferred embodiments may be used and it is intended that the disclosure may be practiced otherwise than as specifically described herein. Accordingly, this disclosure includes all modifications encompassed within the spirit and scope of the disclosure as defined by the claims below.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 38

<210> SEQ ID NO 1
 <211> LENGTH: 29727
 <212> TYPE: DNA
 <213> ORGANISM: Coronavirus
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (265)..(13398)
 <223> OTHER INFORMATION: ORF 1a
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (13398)..(21482)
 <223> OTHER INFORMATION: ORF 1b
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (21492)..(25256)
 <223> OTHER INFORMATION: ORF S
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (25268)..(26089)
 <223> OTHER INFORMATION: ORF X1
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (25689)..(26150)
 <223> OTHER INFORMATION: ORF X2
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (26117)..(26344)
 <223> OTHER INFORMATION: ORF E
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (26398)..(27060)
 <223> OTHER INFORMATION: ORF M
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (27074)..(27262)
 <223> OTHER INFORMATION: ORF X3
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (27273)..(27638)
 <223> OTHER INFORMATION: ORF X4
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (27864)..(28115)
 <223> OTHER INFORMATION: ORF X5
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (28120)..(29385)
 <223> OTHER INFORMATION: ORF N

<400> SEQUENCE: 1

```

ttattagggt tttacctacc caggaaaagc caaccaacct cgatctcttg tagatctggt    60
ctctaaacga actttaaagt ctgtgtagct gtcgctcggc tgcattgcta gtgcacctac    120
gcagataaaa caataataaa ttttactgtc gttgacaaga aacgagtaac tcgtccctct    180
tctgcagact gcttacgggt tcgtccgtgt tgcagtcgat catcagcata cctagggttc    240
gtccgggtgt gaccgaaagg taagatggag agccttggtc ttggtgtcaa cgagaaaaca    300
cacgtccaac tcagtttgcc tgccttcag gttagagacg tgctagtgcg tggcttcggg    360
gactctgtgg aagaggccct atcggaggca cgtgaacacc tcaaaaatgg cacttgtggt    420
ctagtagagc tggaaaagg cgtactgcc cagcttgaac agccctatgt gttcattaaa    480
cgttctgatg ccttaagcac caatcaccgc cacaaggtcg ttgagctggt tgcagaaatg    540
gacggcattc agtacggtcg tagcgggata acactgggag tactcgtgcc acatgtgggc    600
gaaaccccaa ttgataccg caatgttctt cttcgtgaaga acggaataa gggagccggt    660
ggtcatagct atggcatcga tctaaagtct tatgacttag gtgacgagct tggcactgat    720

```

-continued

cccattgaag attatgaaca aaactggaac actaagcatg gcagtggtgc actccgtgaa	780
ctcactcgtg agctcaatgg aggtgcagtc actcgcctatg tcgacaacaa tttctgtggtc	840
ccagatgggt accctcttga ttgcatcaaa gattttctcg cacgcgcggg caagtcaatg	900
tgcactcttt ccgaacaact tgattacatc gagtcgaaga gaggtgtcta ctgctgcccgt	960
gaccatgagc atgaattgc ctgggtcact gagcgcctctg ataagagcta cgagcaccag	1020
acacccttcg aaattaagag tgccaagaaa ttgacactt tcaaagggga atgcccaaag	1080
tttgtgttc ctcttaactc aaaagtcaaa gtcattcaac cacgtgttga aaagaaaaag	1140
actgaggggt tcatggggcg tatacgcctct gtgtaccctg ttgcatctcc acaggagtgt	1200
aacaatatgc acttgtctac cttgatgaaa tgtaatcatt gcgatgaagt ttcattggcag	1260
acgtgcgact ttctgaaagc cacttgtgaa catttgtgca ctgaaaattt agttattgaa	1320
ggacctacta catgtgggta cctacctact aatgctgtag tgaaaatgcc atgtcctgcc	1380
tgtcaagacc cagagattgg acctgagcat agtggtgcag attatcacia ccaactcaaac	1440
attgaaactc gactccgcaa gggaggtagg actagatggt ttggaggctg tgtgtttgcc	1500
tatgttggt gctataataa gcgtgcctac tgggttcctc gtgctagtgc tgatattggtc	1560
tcaggccata ctggcattac tggtgacaat gtggagacct tgaatgagga tctccttgag	1620
atactgagtc gtgaacgtgt taacattaac attgttggcg attttcattt gaatgaagag	1680
gttgccatca ttttggcacc tttctctgct tctacaagtg cctttattga cactataaag	1740
agtcttgatt acaagcttt caaaaccatt gttgagtcct gcgtaacta taaagttacc	1800
aagggaagc ccgtaaaagg tgcttggaac attggacaac agagatcagt ttaaacacca	1860
ctgtgtggtt ttcctcaca ggtgctggt gttatcagat caatttttgc gcgcacactt	1920
gatgcagcaa accactcaat tcttgattg caaagagcag ctgtcccat acttgatggt	1980
atctctgaac agtcattacg tcttgcgac gccatgggtt atactcaga cctgctcacc	2040
aacagtgtca ttattatggc atatgtaact ggtggtcttg tacaacagac ttctcagtg	2100
ttgtctaac ttttggcac tactgtttaa aaactcaggc ctatcttga atggattgag	2160
gcgaaacta gtgcaggagt tgaattctc aaggatgctt gggagattct caaattctc	2220
attacaggtg ttttgacat cgtcaagggt caaatacagg ttgctcaga taacatcaag	2280
gattgtgtaa aatgcttcat tgatgtgtt aacaaggcac tcgaaatgtg cattgatcaa	2340
gtcactatcg ctggcgcaaa gttgcgatca ctcaacttag gtgaagtctt catcgcctca	2400
agcaagggac tttaccgtca gtgtatacgt ggcaaggagc agctgcaact actcatgct	2460
cttaaggcac caaagaagt aaccttctt gaagggtgatt cacatgacac agtacttacc	2520
tctgaggagg ttgttctcaa gaacggtgaa ctogaagcac tcgagacgcc cgttgatagc	2580
ttcaciaaat gagctatcgt tggcacacca gtctgtgtaa atggcctcat gctcttagag	2640
attaaggaca aagaacaata ctgcgcattg tctcctggtt tactggctac aaacaatgtc	2700
tttcgcttaa aaggggtgac accaattaaa ggtgtaacct ttggagaaga tactgtttg	2760
gaagttcaag gttacaagaa tgtgagaatc acatttgagc ttgatgaacg tgttgacaaa	2820
gtgcttaatg aaaagtgtc tgtctacact gttgaatccg gtaccgaagt tactgagttt	2880
gcatgtgttg tagcagaggc tgttgtgaag actttacaac cagtttctga tctccttacc	2940
aacatgggta ttgatcttga tgagtggagt gtagctacat tctacttatt tgatgatgct	3000
ggtgaagaaa acttttcatc acgtatgtat tgttcctttt accctccaga tgaggaagaa	3060

-continued

gaggacgatg	cagagtgtga	ggaagaagaa	attgatgaaa	cctgtgaaca	tgagtacggt	3120
acagaggatg	attatcaagg	tctccctctg	gaatttggtg	cctcagctga	aacagttcga	3180
gttgaggaag	aagaagagga	agactggctg	gatgatacta	ctgagcaatc	agagattgag	3240
ccagaaccag	aacctacacc	tgaagaacca	gttaatcagt	ttactggtta	tttaaaactt	3300
actgacaatg	ttgccattaa	atgtgttgac	atcgtaag	aggcacaag	tgctaactct	3360
atggtgattg	taaatgctgc	taacatacac	ctgaaacatg	gtggtggtgt	agcaggtgca	3420
ctcaacaagg	caaccaatgg	tgccatgcaa	aaggagagtg	atgattacat	taagctaaat	3480
ggccctctta	cagtaggagg	gtcttgtttg	ctttctggac	ataatcttgc	taagaagtgt	3540
ctgcatgttg	ttggacntaa	cctaaaatgca	ggtgaggaca	tccagcttct	taaggcagca	3600
tatgaaaatt	tcaattcaca	ggacatctta	cttgccacct	tgttgtcagc	aggcatatct	3660
ggtgctaaac	cacttcagtc	tttacaagtg	tgcgtagcaga	cggttcgtac	acaggtttat	3720
attgcagtca	atgacaaaagc	tctttatgag	caggttgctca	tggattatct	tgataacctg	3780
aagcctagag	tggaaagcacc	taaacaagag	gagccaccaa	acacagaaga	ttccaaaact	3840
gaggagaaat	ctgtcgtaca	gaagcctgtc	gatgtgaagc	caaaaattaa	ggcctgcatt	3900
gatgaggtta	ccacaacact	ggaagaaact	aagtttctta	ccaataagtt	actcttgttt	3960
gctgatatca	atggttaagct	ttaccatgat	tctcagaaca	tgcttagagg	tgaagatatg	4020
tctttccttg	agaaggatgc	accttacatg	gtaggtgatg	ttatcactag	tggtgatatc	4080
acttggttg	taataccttc	caaaaaggct	ggtggcacta	ctgagatgct	ctcaagagct	4140
ttgaagaaag	tgccagttga	tgagtatata	accacgtacc	ctggacaagg	atgtgctggt	4200
tatacacttg	aggaagctaa	gactgctctt	aagaaatgca	aatctgcatt	ttatgtacta	4260
ccttcagaag	cacctaagtc	taaggaagag	attctaggaa	ctgtatcctg	gaatttgaga	4320
gaaatgcttg	ctcatgctga	agagacaaga	aaattaatgc	ctatatgcat	ggatgttaga	4380
gccataatgg	caaccatcca	acgtaagtat	aaaggaatta	aaattcaaga	gggcatcggt	4440
gactatggtg	tccgattcct	cttttatact	agtaagagc	ctgtagcttc	tattattacg	4500
aagctgaact	ctctaataatga	gccgcttgtc	acaatgcca	ttggttatgt	gacacatggt	4560
tttaactctg	aagaggetgc	gcgctgtatg	cgttctctta	aagctcctgc	cgtagtgtca	4620
gtatcatcac	cagatgctgt	tactacatat	aatggatacc	tcacttcgtc	atcaaagaca	4680
tctgaggagc	actttgtaga	aacagtttct	ttggctggct	cttacagaga	ttggctctat	4740
tcaggacagc	gtacagagtt	aggtgttgaa	tttcttaagc	gtggtgacaa	aattgtgtac	4800
cacactctgg	agagccccgt	cgagtttcat	cttgacggtg	aggttcttcc	acttgacaaa	4860
ctaaagagtc	tcttatccct	gcgggaggtt	aagactataa	aagtgttcac	aactgtggac	4920
aacactaatc	tccacacaca	gcttggtgat	atgtctatga	catatggaca	gcagtttggt	4980
ccaacatact	tggatggtgc	tgatgttaca	aaaattaaac	ctcatgtaaa	tcatgaggg	5040
aagactttct	ttgtactacc	tagtgatgac	acactacgta	gtgaagcttt	cgagtactac	5100
catactcttg	atgagagttt	tcttggtagg	tacatgtctg	ctttaaacca	cacaaagaaa	5160
tggaaatttc	ctcaagttgg	tggtttaact	tcaattaaat	gggctgataa	caattgttat	5220
ttgtctagtg	ttttattagc	acttcaacag	cttgaagtca	aattcaatgc	accagcactt	5280
caagaggctt	attatagagc	cctgctggtg	gatgctgcta	acttttgctc	actcactctc	5340
gcttacagta	ataaaactgt	tggcgagctt	ggtgatgtca	gagaaactat	gacctactct	5400
ctacagcatg	ctaatttggg	atctgcaaaag	cgagttctta	atgtggtgtg	taaacattgt	5460

-continued

ggtcagaaaa ctactacctt aacgggtgta gaagctgtga tgtatatggg tactctatct	5520
tatgataatc ttaagacagg tgtttccatt ccatgtgtgt gtggtcgtga tgctacacaa	5580
tatctagtagt aacaagagtc ttcttttgggt atgatgtctg caccacctgc tgagtataaa	5640
ttacagcaag gtacattcct atgtgcgaat gagtacctg gtaactatca gtgtggctcat	5700
tacactcata taactgctaa ggagaccctc tatcgtattg acggagctca ccttacaag	5760
atgtcagagt acaaaggacc agtgactgat gttttctaca aggaacatc ttacactaca	5820
accatcaagc ctgtgtcgta taaactcgat ggagttactt acacagagat tgaacaaaa	5880
ttggatgggt attataaaaa ggataatgct tactatacag agcagcctat agaccttgta	5940
ccaactcaac cattacaaa tgcgagtttt gataatttca aactcactg ttctaacaca	6000
aaatttgctg atgatttaaa tcaaatgaca ggcttcacaa agccagcttc acgagagcta	6060
tctgtccatc tcttccaga cttgaatggc gatgtagtgg ctattgacta tagacactat	6120
tcagcgagtt tcaagaaagg tgctaaatta ctgcataagc caattgtttg gcacattaac	6180
caggctacaa ccaagacaac gttcaaacca aacacttggt gtttacgttg tctttggagt	6240
acaaagccag tagatacttc aaattcattt gaagttctgg cagtagaaga cacacaagga	6300
atggacaatc ttgcttgtga aagtcaacaa cccacctctg aagaagtagt ggaaatcct	6360
accatacaga aggaagtcac agagtgtgac gtgaaaacta ccgaagttgt aggcaatgac	6420
atacttaaac catcagatga aggtgttaaa gtaacacaag agttaggcca tgaggatcct	6480
atggctgctt atgtggaaaa cacaagcatt accattaaga aacctaatga gctttcacta	6540
gccttagggt taaaaacaat tgcacctcat ggtattgctg caattaatag tgttccttgg	6600
agtaaaattt tggcttatgt caaaccttc ttaggacaag cagcaattac aacatcaaat	6660
tgcgctaaga gattagcaca acgtgtgttt aacaattata tgccttatgt gtttacatta	6720
ttgttccaat tgtgtacttt tactaaaagt accaattcta gaattagagc ttactacct	6780
acaactattg ctaaaaatag tgttaagagt gttgctaact tatgtttgga tgccggcatt	6840
aattatgtga agtcacccaa attttctaaa ttgttcacaa tcgctatgtg gctattgttg	6900
ttaagtattt gcttaggttc tctaactctgt gtaactgctg cttttggtgt actcttatct	6960
aattttggtg ctcttctta ttgtaatggc gttagagaat tgatcttaa tctgtctaac	7020
gttactacta tggatttctg tgaaggttct tttccttgca gcatttgttt aagtggatta	7080
gactcccttg attcttatcc agctcttgaa accattcagg tgacgatttc atcgtaacag	7140
ctagacttga caattttagg tctggccgct gagtgggttt tggcatatat gttgttcaca	7200
aaattctttt atttattagg tctttcagct ataatgcagg tgttctttgg ctattttgct	7260
agtcatttca tcagcaatc ttggctcatg tggtttatca ttagtattgt acaaatggca	7320
cccgtttctg caatggttag gatgtacac ttctttgctt ctttctacta catatggaag	7380
agctatgttc atatcatgga tggttgcacc tcttcgactt gcatgatgtg ctataagcgc	7440
aatcgtgcc aacgcgttga gtgtacaact attgtaatg gcatgaagag atctttctat	7500
gtctatgcaa atggaggccg tggcttctgc aagactcaca attggaattg tctcaattgt	7560
gacacatttt gcactggtag tacattcatt agtgaatgaag ttgctcgtga tttgtcactc	7620
cagtttaaaa gaccaatcaa cctactgac cagtcacgt atattgttga tagtgttgc	7680
gtgaaaaatg ggcgcttca cctctacttt gacaaggctg gtcaaaagac ctatgagaga	7740
catccgctct cccattttgt caatttagac aatttgagag ctaacaacac taaaggttca	7800

-continued

ctgcctatta atgtcatagt ttttgatggc aagtccaaat ggcagcagtc tgcttctaag	7860
tctgcttctg tgtactacag tcagctgatg tgccaaccta ttctggttgc tgaccaagtt	7920
cttgatcag acgttgaga tagtactgaa gtttccgta agatgttga tgcttatgtc	7980
gacacctttt cagcaacttt tagtgttcct atggaaaaac ttaaggcact tgttgctaca	8040
gctcacagcg agttagcaaa ggggttagct ttagatggtg tctttctac attcgtgtca	8100
gctgcccgac aagggtgtgt tgataccgat gttgacacaa aggatgttat tgaatgtctc	8160
aaactttcac atcactctga cttagaagtg acaggtgaca gttgtaaca tttcatgctc	8220
acctataata aggttgaana catgacgcc agagatcttg ggcgatgtat tgactgtaat	8280
gcaaggcata tcaatgccca agtagcaaaa agtcacaatg tttcactcat ctggaatgta	8340
aaagactaca tgtctttatc tgaacagctg cgtaaacaaa ttctgtagtgc tgccaagaag	8400
aacaacatac cttttagact aacttgtgct acaactagac aggttgtcaa tgtcataact	8460
actaaaatct cactcaaggg tggttaagatt gttagtactt gttttaact tatgcttaag	8520
gccacattat tgtgcttct tgctgcattg gtttgttata tctttatgcc agtacatata	8580
ttgtcaatcc atgatggta cacaaatgaa atcattggtt acaaagccat tcaggatggt	8640
gtcactcgtg acatcatttc tactgatgat tgttttgcaa ataaacatgc tggttttgac	8700
gcatggttta gccagcgtgg tggttcatac aaaaatgaca aaagctgcc tgtagtagct	8760
gctatcatta caagagagat tggtttcata gtgcctggct taccgggtac tgtgctgaga	8820
gcaatcaatg gtgacttctt gcattttcta cctcgtggtt ttagtgctgt tggcaacatt	8880
tgctacacac cttccaaact cattgagat agtgattttg ctacctctgc ttgcttctt	8940
gctgctgagt gtacaatttt taaggatgct atgggcaaac ctgtgccata ttgttatgac	9000
actaatttgc tagagggttc tatttcttat agtgagcttc gtccagacac tcttatggtg	9060
cttatggatg gttccatcat acagtttctt aacacttacc tggagggttc tgttagagta	9120
gtaacaactt ttgatgctga gtactgtaga catggtacat gcgaaaggct agaagtaggt	9180
atttgcctat ctaccagtg tagatgggtt ctttaaatg agcattacag agctctatca	9240
ggagttttct gtggtgttga tgcgatgaat ctcatagcta acatctttac tctcttctg	9300
caacctgtgg gtgctttaga tgtgtctgct tcagtagtgg ctggtggtat tattgccata	9360
ttggtgactt gtgctgcta ctactttatg aaattcagac gtgtttttgg tgagtacaac	9420
catgttgttg ctgctaagc acttttgtt ttgatgtctt tcactatact ctgtctggtg	9480
ccagcttaca gctttctgcc gggagctcac tcagctttt acttgactt gacattctat	9540
ttcaccaatg atgtttcatt cttggctcac cttcaatggt ttgccatggt ttctctatt	9600
gtgcctttt ggataacagc aatctatgta ttctgtattt ctctgaagca ctgccattgg	9660
ttctttaaca actatcttag gaaaagagtc atgtttaatg gagttacatt tagtacctc	9720
gaggaggctg ctttgtgtac cttttgtctc aacaaggaaa tgtacctaaa attgcgtagc	9780
gagacactgt tgccacttac acagtataac aggtatcttg ctctatataa caagtacaag	9840
tatttcagtg gagccttaga tactaccagc tatcgtgaag cagcttgctg ccacttagca	9900
aaggctctaa atgactttag caactcaggt gctgatgttc tctaccaacc accacagaca	9960
tcaatcactt ctgctgttct gcagagtggt tttaggaaaa tggcattccc gtcaggcaaa	10020
gttgaagggt gcatggtaca agtaacctgt ggaactacaa ctcttaatgg attgtggttg	10080
gatgacacag tatactgtcc aagacatgct atttgcacag cagaagacat gcttaacct	10140
aaactatgaag atctgctcat tcgcaaatcc aaccatagct ttcttgttca ggctggcaat	10200

-continued

gttcaacttc gtgttattgg ccattctatg caaaattgtc tgcttaggct taaagttgat 10260
 acttctaacc ctaagacacc caagtataaa tttgtccgta tccaacctgg tcaaacattt 10320
 tcagttctag catgctacaa tggttcacca tctggtggtt atcagtggtc catgagacct 10380
 aatcatacca ttaaaggttc tttccttaat ggatcatgtg gtagtggtgg ttttaacatt 10440
 gattatgatt gcgtgtcttt ctgctatatg catcatatgg agcttccaac aggagtacac 10500
 gctggtactg acttagaagg taaattctat ggtccatttg ttgacagaca aactgcacag 10560
 gctgcaggta cagacacaac cataacatta aatgttttgg catggctgta tgctgctgtt 10620
 atcaatggty atagggtggt tcttaataga ttcaccacta ctttgaatga ctttaacctt 10680
 gtggcaatga agtacaacta tgaacctttg acacaagatc atgttgacat attgggacct 10740
 ctttctgctc aaacaggaat tgcgctctta gatatgtgtg ctgctttgaa agagctgctg 10800
 cagaatggta tgaatggtcg tactatcctt ggtagcacta ttttagaaga tgagtttaca 10860
 ccatttgatg ttgtagaca atgctctggt gttaccttc aaggttaagtt caagaaaatt 10920
 gtttaaggca ctcactatg gatgctttta actttcttga catcactatt gattcttgtt 10980
 caaagtacac agtggctact gttttcttt gtttacgaga atgctttctt gccatttact 11040
 cttggtatta tggcaattgc tgcatgtgct atgctgcttg ttaagcataa gcaacgattc 11100
 ttgtgcttgt ttctgttacc ttctcttga acagttgctt actttaatat ggtctacatg 11160
 cctgctagct ggggatgctg tatcatgaca tggcttgaat tggctgacac tagcttgtct 11220
 ggttataggc ttaaggattg tgttatgtat gcttcagctt tagttttgct tattctcatg 11280
 acagctcgca ctgtttatga tgatgctgct agacgtgttt ggacactgat gaatgtcatt 11340
 acacttgttt acaaagtcta ctatggtaat gcttttagatc aagctatttc catgtgggce 11400
 ttagttatct ctgtaacctc taactattct ggtgtcgtta cgactatcat gtttttagct 11460
 agagctatag tgtttgtgtg tgttgagtat taccattgt tatttattac tggcaacacc 11520
 ttacagtgtg tcatgcttgt ttattgttct ttaggctatt gttgctgctg ctactttggc 11580
 ctttctgtt tactcaaccg ttacttcagg cttactcttg gtgtttatga ctacttggc 11640
 tctacacaag aatttaggta tatgaacctc caggggcttt tgcctcctaa gagtagtatt 11700
 gatgctttca agcttaacat taagtgttg ggtattggag gtaaaccatg tatcaagggt 11760
 gctactgtac agtctaaaat gtctgacgta aagtgcacat ctgtggtact gctctcgggt 11820
 cttcaacaac ttagagtaga gtcacttctt aaattgtggg cacaatgtgt acaactccac 11880
 aatgatattc ttcttgcaaa agacacaact gaagctttcg agaagatggt ttctcttttg 11940
 tctgttttgc tatccatgca ggggtgctga gacattaata ggttggtcga ggaaatgctc 12000
 gataaccgtg ctactcttca ggetattgct tcagaattta gttctttacc atcatatgcc 12060
 gcttatgcc a ctgcccagga ggcctatgag caggctgtag ctaatggtga ttctgaagtc 12120
 gttctcaaaa agttaaagaa atctttgaat gtggctaaat ctgagtttga ccgtgatgct 12180
 gccatgcaac gcaagttgga aaagatggca gatcaggcta tgaccocaaat gtacaaacag 12240
 gcaagatctg aggacaagag ggcaaaagta actagtgtca tgcaacaat gctcttctact 12300
 atgcttagga agcttgataa tgatgcactt aacaacatta tcaacaatgc gcgtgatggt 12360
 tgtgttccac tcaacatcat accattgact acagcagcca aactcatggt tgttgtccct 12420
 gattatggta cctacaagaa cacttgtgat ggtaaacact ttacatatgc atctgcactc 12480
 tgggaaatcc agcaagttgt tgatgcggat agcaagattg ttcaacttag tgaaattaac 12540

-continued

atggacaatt	caccaaattt	ggettggcct	cttattgtta	cagctctaag	agccaactca	12600
gctgttaaac	tacagaataa	tgaactgagt	ccagtagcac	tacgacagat	gtcctgtgcg	12660
gctggtacca	cacaaacagc	ttgtactgat	gacaatgcac	ttgcctacta	taacaattcg	12720
aagggaggta	ggtttgtgct	ggcattacta	tcagaccacc	aagatctcaa	atgggctaga	12780
ttccctaaga	gtgatggtac	aggtacaatt	tacacagaac	tggaaccacc	ttgtaggttt	12840
gttacagaca	cacaaaagg	gcctaaagtg	aaatacttgt	acttcatcaa	aggcttaaac	12900
aacctaaata	gaggtatggt	gctgggcagt	ttagctgcta	cagtacgtct	tcaggctgga	12960
aatgctacag	aagtacctgc	caattcaact	gtgctttcct	tctgtgcttt	tgcagtagac	13020
cctgctaaag	catataagga	ttacctagca	agtggaggac	aaccaatcac	caactgtgtg	13080
aagatgttgt	gtacacacac	tggtacagga	caggcaatta	ctgtaacacc	agaagctaac	13140
atggaccaag	agtcccttgg	tggtgcttca	tgttgtctgt	attgtagatg	ccacattgac	13200
catccaaatc	ctaaggatt	ctgtgacttg	aaaggtaaat	acgtccaaat	acctaccact	13260
tgtgctaatt	accagtgagg	ttttacactt	agaaacacag	tctgtaccgt	ctgcggaatg	13320
tggaaggtt	atggctgtag	ttgtgaccaa	ctccgcgaac	ccttgatgca	gtctgcggat	13380
gcatcaacgt	ttttaaaccg	gtttgcggtg	taagtgcagc	ccgtcttaca	ccgtgcgcca	13440
caggcactag	tactgatgtc	gtctacaggg	cttttgatat	ttacaacgaa	aaagttgctg	13500
gttttgcaaa	gttcctaaaa	actaattgct	gtcgcttcca	ggagaaggat	gaggaaggca	13560
atattattaga	ctcttacttt	gtagttaaga	ggcatactat	gtctaactac	caacatgaag	13620
agactattta	taacttggtt	aaagattgtc	cagcggttgc	tgtccatgac	tttttcaagt	13680
ttagagtaga	tggtgacatg	gtaccacata	tatcacgtca	gcgtctaact	aaatacacia	13740
tggctgattt	agtctatgct	ctacgtcatt	ttgatgaggg	taattgtgat	acattaaaag	13800
aaatactcgt	cacatacaat	tgtctgtgat	atgattattt	caataagaag	gattgggatg	13860
acttcgtaga	gaatcctgac	atcttacgcg	tatatgctaa	cttaggtgag	cgtgtacgcc	13920
aatcattatt	aaagactgta	caattctgcg	atgctatgcg	tgatgcaggc	attgtaggcg	13980
tactgacatt	agataatcag	gatcttaatg	ggaactggta	cgatttcggt	gatttcgtac	14040
aagtagcacc	aggctgcgga	gttcctattg	tggattcata	ttactcattg	ctgatgccca	14100
tctcactttt	gactagggca	ttggctgctg	agtcccatat	ggatgctgat	ctcgcaaaac	14160
cacttattaa	gtgggatttg	ctgaaatatg	attttacgga	agagagactt	tgtctcttcg	14220
accgttattt	taaatattgg	gaccagacat	accatcccaa	ttgtattaac	tgtttggtatg	14280
ataggtgat	ccttcattgt	gcaaacctta	atgtgttatt	ttctactgtg	tttccacctt	14340
caagttttgg	accactagta	agaaaaatat	ttgtagatgg	tgttcctttt	gttgtttcaa	14400
ctggatacca	ttttcgtgag	ttaggagtcg	tacataatca	ggatgtaaac	ttacatagct	14460
cgcgtctcag	tttcaaggaa	cttttagtgt	atgctgctga	tccagctatg	catgcagctt	14520
ctggcaattt	attgctagat	aaacgcacta	catgcttttc	agtagctgca	ctaacaacaa	14580
atggtgcttt	tcaaactgtc	aaacccggtt	attttaataa	agacttttat	gactttgctg	14640
tgtctaaagg	tttctttaag	gaaggaagtt	ctggtgaaat	aaaacacttc	ttctttgctc	14700
aggatggcaa	cgctgctatc	agtgattatg	actattatcg	ttataatctg	ccaacaatgt	14760
gtgatatcag	acaactccta	tctgtagttg	aagttgttga	taaatacttt	gattgttacg	14820
atggtggctg	tattaatgcc	aaccaagtaa	tctgtaacaa	tctggataaa	tcagctgggt	14880
tcccatttaa	taaatggggg	aaggctagac	tttattatga	ctcaatgagt	tatgaggatc	14940

-continued

```

aagatgcact tttcgcgtat actaagcgtat atgtcatccc tactataact caaatgaatc 15000
ttaagtatgc cattagtgca aagaatagag ctgcgaccgt agctgggtgc tctatctgta 15060
gtactatgac aaatagacag ttcatcaga aattattgaa gtcaatagcc gccactagag 15120
gagctactgt ggtaattgga acaagcaagt tttacgggtg ctggcataat atgttaaaaa 15180
ctgtttacag tgatgtagaa actccacacc ttatgggttg ggattatcca aaatgtgaca 15240
gagccatgcc taacatgctt aggataatgg cctctcttgt tcttgctcgc aaacataaca 15300
cttgctgtaa cttatcacac cgtttctaca ggtagctaa cgagtgtgcg caagtattaa 15360
gtgagatggt catgtgtggc ggctcactat atgttaaac aggtggaaca tcatccggtg 15420
atgctacaac tgcttatgct aatagtgtct ttaacatttg tcaagctgtt acagccaatg 15480
taaatgcact tctttcaact gatggtaata agatagctga caagtatgct cgcaatctac 15540
aacacaggct ctatgagtgt ctctatagaa atagggatgt tgatcatgaa ttcgtggatg 15600
agttttacgc ttacctgcgt aaacatttct ccatgatgat tctttctgat gatgccggtg 15660
tgtgctataa cagtaactat gcggctcaag gtttagtagc tagcattaag aactttaagg 15720
cagttcttta ttatcaaaat aatgtgttca tgtctgagggc aaaatgttgg actgagactg 15780
accttactaa aggacctcac gaattttgct cacagcatac aatgctagtt aaacaaggag 15840
atgattacgt gtacctgctt taccagatc catcaagaat attagggcga ggctgttttg 15900
tcgatgatat tgtcaaaaca gatggtacac ttatgattga aaggttcgtg tcaactggcta 15960
ttgatgctta cccacttaca aaacatccta atcaggagta tgctgatgct tttcaactgt 16020
atttacaata cattagaaag ttacatgatg agcttactgg ccacatgttg gacatgtatt 16080
ccgtaatgct aactaatgat aacacctcac ggtactggga acctgagttt tatgaggcta 16140
tgtacacacc acatacagtc ttgcaggctg taggtgcttg tgtattgtgc aattcacaga 16200
cttcacttcg ttgcgggtcc tgtattagga gaccattcct atgttgcaag tgctgctatg 16260
accatgtcat ttcaacatca cacaaattag tgttgtctgt taatccctat gtttgcaatg 16320
cccagggtg tgatgtcact gatgtgacac aactgtatct aggaggtatg agctattatt 16380
gcaagtcaaa taagcctccc attagtttcc cattatgtgc taatggtcag gtttttggtt 16440
tatacaaaaa cacatgtgta ggcagtgaca atgtcactga cttcaatgag atagcaacat 16500
gtgattggac taatgctggc gattacatac ttgccaacac ttgtactgag agactcaagc 16560
ttttcgagc agaaacgctc aaagccactg aggaaacatt taagctgtca tatggtattg 16620
ctactgtacg cgaagtactc tctgacagag aattgcatct ttcattggag gttgaaaaac 16680
ctagaccacc attgaacaga aactatgtct ttactgggta ccgtgtaact aaaaatagta 16740
aagtacagat tggagagtac acctttgaaa aaggtgacta tgggtgatgct gttgtgtaca 16800
gaggtactac gacatacaag ttgaatgttg gtgattactt tgtgttgaca tctcacactg 16860
taatgccact tagtgacctc actctagtgc cacaaagca ctatgtgaga attactggct 16920
tgtaccaaac actcaacatc tcagatgagt tttctagcaa tggttgcaat tatcaaaagg 16980
tcggcatgca aaagtactct aactccaag gaccacctgg tactggtaag agtcattttg 17040
ccateggact tgctctctat taccatctg ctgcgatagt gtatcggca tgctctcatg 17100
cagctgttga tgccctatgt gaaaaggcat taaaatattt gcccatagat aaatgtagta 17160
gaatcatacc tgcgctgctg cgcgtagagt gttttgataa attcaaagtg aattcaacac 17220
tagaacagta tgtttctgct actgtaaagt cattgccaga aacaactgct gacattgtag 17280

```

-continued

tctttgatga	aatctctatg	gctactaatt	atgacttgag	tgttgtcaat	gctagacttc	17340
gtgcaaaaca	ctacgtctat	attggcgatc	ctgctcaatt	accagcccc	cgcacattgc	17400
tgactaaagg	cacactagaa	ccagaatatt	ttaattcagt	gtgcagactt	atgaaaaaca	17460
taggtocaga	catgttcctt	ggaacttgc	gccgttgtcc	tgctgaaatt	gttgacactg	17520
tgagtgcttt	agtttatgac	aataagctaa	aagcacacaa	ggataagtca	gctcaatgct	17580
tcaaaatggt	ctacaaaggt	gttattacac	atgatgttcc	atctgcaatc	aacagacctc	17640
aaataggcgt	tgtaagagaa	tttcttacac	gcaatcctgc	ttggagaaaa	gctgttttta	17700
tctcacctta	taattcacag	aacgctgtag	cttcaaaaat	cttaggattg	cctacgcaga	17760
ctggtgattc	atcacagggt	tctgaatatg	actatgtcat	attcacacaa	actactgaaa	17820
cagcacactc	ttgtaatgtc	aaccgcttca	atgtggctat	cacaagggca	aaaattggca	17880
ttttgtgcat	aatgtctgat	agagatcttt	atgacaaact	gcaatttaca	agtctagaaa	17940
taccacgtcg	caatgtggct	acattacaag	cagaaaatgt	aactggactt	tttaaggact	18000
gtagtaagat	cattactggt	cttcatccta	cacaggcacc	tacacacctc	agcgttgata	18060
taaagttoaa	gactgaagga	ttatgtgttg	acataccagg	cataccaaag	gacatgacct	18120
accgtagact	catctctatg	atgggtttca	aaatgaatta	ccaagtcaat	ggttacccta	18180
atatgtttat	caccocgcaa	gaagctatc	gtcacgttcg	tgcggtgatt	ggctttgatg	18240
tagagggctg	tcatgcaact	agagatgctg	tggtactaa	cctacctctc	cagctaggat	18300
tttctacagg	tgtaactta	gtagctgtac	cgactggta	tgttgacact	gaaaataaca	18360
cagaattcac	cagagttaat	gcaaacctc	caccagggtga	ccagtttaaa	catcttatac	18420
cactcatgta	taaaggcttg	ccttgaatg	tagtgcgtat	taagatagta	caaatgctca	18480
gtgatacact	gaaaggattg	tcagacagag	tcgtgttcgt	cctttggcg	catggctttg	18540
agcttacatc	aatgaagtac	tttgtcaaga	ttggacctga	aagaacgtgt	tgtctgtgtg	18600
acaaacgtgc	aacttgcttt	tctacttcat	cagatactta	tgcttctgg	aatcattctg	18660
tgggttttga	ctatgtctat	aaccattta	tgattgatgt	tcagcagtgg	ggctttacgg	18720
gtaaccttca	gagtaacct	gaccaacatt	gccaggta	tggaaatgca	catgtggcta	18780
gttgtgatgc	tatcatgact	agatgtttag	cagtcctga	gtgctttgtt	aagcgcgttg	18840
attggtctgt	tgaataacct	attataggag	atgaactgag	ggtaattct	gcttgacaga	18900
aagtacaaca	catggttggt	aagtctgcat	tgcttctga	taagtttcca	gttcttcatg	18960
acattgga	tccaaaggct	atcaagtgtg	tgctcaggc	tgaagtagaa	tggaagtctt	19020
acgatgctca	gccatgtagt	gacaaagctt	acaaaataga	ggagctcttc	tattcttatg	19080
ctacacatca	cgataaattc	actgatggtg	tttgtttgtt	ttggaattgt	aacgttgatc	19140
gttaccacgc	caatgcaatt	gtgtgtaggt	ttgacacaag	agccttgta	aacttgaact	19200
taccaggctg	tgatgggtgt	agtttgtatg	tgaataagca	tgattccac	actccagctt	19260
tcgataaaag	tgcatctact	aatttaaagc	aattgccttt	cttttactat	tctgatagtc	19320
cttgtgagtc	tcattggcaa	caagtagtgt	cggatattga	ttatgttcca	ctcaaatctg	19380
ctacgtgtat	tacacgatgc	aatttaggtg	gtgctgtttg	cagacacat	gcaaatgagt	19440
accgacagta	cttggatgca	tataaatatga	tgatttctgc	tggatttagc	ctatggattt	19500
acaaacaatt	tgatacttat	aacctgtgga	atacatttac	caggttacag	agtttagaaa	19560
atgtggctta	taatgttgtt	aataaaggac	actttgatgg	acacgccggc	gaagcacctg	19620
tttccatcat	taataatgct	gtttacacaa	aggtagatgg	tattgatgtg	gagatctttg	19680

-continued

aaaataagac aacacttccct gttaatggtg catttgagct ttgggctaag cgtaacatta 19740
 aaccagtgcc agagattaag atactcaata atttgggtgt tgatatacgt gctaatactg 19800
 taatctggga ctacaaaaga gaagccccag cacatgtatc tacaataggt gtctgcacaa 19860
 tgactgacat tgccaagaaa cctactgaga gtgcttggtc ttcacttact gtcttgtttg 19920
 atggtagagt ggaaggacag gttagacctt ttagaaacgc ccgtaatggt gttttaataa 19980
 cagaaggttc agtcaaaggc ctaaacctt caaagggacc agcacaagct agcgtcaatg 20040
 gagtcacatt aattggagaa tcagtaaaaa cacagtttaa ctactttaag aaagtagacg 20100
 gcattattca acagttgctt gaaacctact ttactcagag cagagactta gaggatttta 20160
 agcccagatc acaaatggaa actgactttc tcgagctcgc tatggatgaa ttcatacagc 20220
 gatataagct cgagggtat gccttcgaac acatcgttta tggagatttc agtcatggac 20280
 aacttgccgg tcttcattta atgataggct tagccaagcg ttcacaagat tcaccactta 20340
 aattagagga ttttatccct atggacagca cagtgaaaaa ttacttcata acagatgcgc 20400
 aaacaggttc atcaaatgt gtgtgttctg tgattgatct tttacttgat gactttgtcg 20460
 agataataaa gtcacaagat ttgtcagtga tttcaaaagt ggtcaaggtt acaattgact 20520
 atgctgaaat ttcattcatg ctttgggtga aggatggaca tggtgaaacc ttctacccaa 20580
 aactacaagc aagcaagcg tggcaaccag gtggtgcgat gcctaactg tacaagatgc 20640
 aaagaatgct tcttgaagg tgtgacctc agaattatgg tgaaaatgct gttataccaa 20700
 aaggaataat gatgaatgtc gcaaagtata ctcaactgtg tcaatactta aatacactta 20760
 ctttagctgt accctacaac atgagagtta ttcactttgg tgctggctct gataaaggag 20820
 ttgcaccagg tacagctgtg ctacagacaat ggttgccaac tggcacacta cttgtcgatt 20880
 cagatcttaa tgacttcgtc tccgacgcag attctacttt aattggagac tgtgcaacag 20940
 tacatacggc taataaatgg gaccttatta ttagegatat gtatgacctt aggaccaaac 21000
 atgtgacaaa agagaatgac tctaaagaag ggtttttcac ttatctgtgt ggatttataa 21060
 agcaaaaact agccctgggt ggttctatag ctgtaaagat aacagagcat tcttggaatg 21120
 ctgaccttta caagcttatg ggccatttct catggtggac agcttttgtt acaaatgtaa 21180
 atgcatcatc atcgggaagca tttttaattg gggctaacta tcttggcaag ccgaaggaac 21240
 aaattgatgg ctataccatg catgctaact acattttctg gaggaacaca aatcctatcc 21300
 agttgtcttc ctattcactc tttgacatga gcaaatttcc tcttaaatga agaggaactg 21360
 ctgtaatgct tcttaaggag aatcaaatca atgatatgat ttattctctt ctggaaaaag 21420
 gtaggcttat cattagagaa aacaacagag ttgtggttcc aagtgatatt cttgttaaca 21480
 actaaacgaa catgtttatt ttcttattat ttcttactct cactagtggg agtgacctg 21540
 accggtgcac cacttttgat gatgttcaag ctccataata cactcaacat acttcatcta 21600
 tgaggggggt ttactatcct gatgaaattt ttagatcaga cactctttat ttaactcagg 21660
 atttatttct tccattttat tctaattgta cagggtttca tactattaat catacgtttg 21720
 gcaacctgt catacctttt aaggatggtt tttattttgc tgccacagag aatcaaatg 21780
 ttgtccgtgg ttgggttttt ggttctacca tgaacaacaa gtcacagtcg gtgattatta 21840
 ttaacaattc tactaatggt gttatacagag catgtaactt tgaattgtgt gacaacctt 21900
 tctttgctgt ttctaaaccc atgggtacac agacacatac tatgatattc gataatgcat 21960
 ttaattgcac tttcgagtac atatctgatg ccttttcgct tgatgtttca gaaaagtacg 22020

-continued

gtaattttaa	acacttacga	gagtttgtgt	ttaaaaaata	agatggggtt	ctctatgttt	22080
ataagggcta	tcaacctata	gatgtagttc	gtgatctacc	ttctgggttt	aacactttga	22140
aacctatfff	taagttgctt	cttggtatta	acattacaaa	ttttagagcc	attcttacag	22200
ccttttcacc	tgctcaagac	atttggggca	cgtcagctgc	agcctatfff	gttggetatt	22260
taaagccaac	tacatttatg	ctcaagtatg	atgaaaatgg	tacaatcaca	gatgctggtg	22320
attgtttca	aaatccactt	gctgaactca	aatgctctgt	taagagcttt	gagattgaca	22380
aaggaattta	ccagacctct	aatttcaggg	ttgttccctc	aggagatggt	gtgagattcc	22440
ctaataattac	aaacttgtgt	ccttttgag	aggtttttaa	tgctactaaa	ttcccttctg	22500
tctatgcatg	ggagagaaaa	aaaatttcta	attgtgttgc	tgattactct	gtgctctaca	22560
actcaacatt	tttttcaacc	tttaagtgt	atggcgtttc	tgccactaag	ttgaatgatc	22620
tttgcttctc	caatgtctat	gcagattctt	ttgtagtcaa	gggagatgat	gtaagacaaa	22680
tagcgccagg	acaaactggg	gttattgctg	attataatta	taaattgcca	gatgatttca	22740
tgggttgtgt	ccttgcttgg	aatactagga	acattgatgc	tacttcaact	ggtaattata	22800
attataaata	taggtatctt	agacatggca	agcttaggcc	ctttgagaga	gacatatcta	22860
atgtgccttt	ctcccctgat	ggcaaacctt	gcaccccacc	tgctcttaat	tgttattggc	22920
cattaatga	ttatggtttt	tacaccacta	ctggcattgg	ctaccaacct	tacagagttg	22980
tagtactttc	ttttgaactt	ttaaatgcac	cggccacggg	ttgtggacca	aaattatcca	23040
ctgaccttat	taagaaccag	tgtgtcaatt	ttaattttaa	tggactcact	ggtactgggtg	23100
tgtaactcc	ttcttcaag	agatttcaac	catttcaaca	atrtggccgt	gatgtttctg	23160
atrtcaactga	ttccgttcca	gatcctaaaa	catctgaaat	attagacatt	tcaccttgct	23220
crtttggggg	tgtaagtgt	attacacctg	gaacaaatgc	ttcatctgaa	gttgctgttc	23280
tatatcaaga	tgtaactgca	actgatgttt	ctacagcaat	tcatgcagat	caactcacac	23340
cagcttggcg	catatattct	actggaaaca	atgtattcca	gactcaagca	ggctgtctta	23400
taggagctga	gcattgtgac	acttcttatg	agtgcgacat	tcctattgga	gctggcattt	23460
gtgctagtta	ccatacagtt	tctttattac	gtagtactag	ccaaaaatct	attgtggctt	23520
atactatgtc	tttaggtgct	gatagttaa	ttgcttactc	taataacacc	attgctatac	23580
ctactaactt	ttcaattagc	attactacag	aagtaatgcc	tgtttctatg	gctaaaaacct	23640
ccgtagattg	taatatgtac	atctgcggag	attctactga	atgtgctaat	ttgcttctcc	23700
aatatggtag	crtttgcaca	caactaaatc	gtgcactctc	aggtattgct	gctgaacagg	23760
atcgcaacac	acgtgaagtg	ttcgctcaag	tcaaacaaat	gtacaaaacc	ccaactttga	23820
aatatrttgg	tggtrttaat	ttttcacaaa	tattacctga	ccctctaaag	ccaactaaga	23880
ggctctttat	tgaggacttg	ctctttaata	aggtagacct	cgctgatgct	ggcttcatga	23940
agcaatatgg	cgaatgccta	ggtgatatta	atgctagaga	tctcatttgt	gctgagaagt	24000
tcaatggact	tacagtgttg	ccacctctgc	tactgatga	tatgattgct	gcctacactg	24060
ctgctctagt	tagtggtagt	gccactgctg	gatggacatt	tggtagctggc	gctgctcttc	24120
aaataccttt	tgctatgcaa	atggcatata	ggttcaatgg	cattggagtt	accctaaatg	24180
ttctctatga	gaacccaaaa	caaatcgcca	accaatttaa	caaggcgatt	agtcaaatc	24240
aagaatcact	tacaacaaca	tcaactgcat	tgggcaagct	gcaagacgtt	gttaaccaga	24300
atgctcaagc	attaacacac	crttgtaaac	aacttagctc	taatrttggg	gcaatrtcaa	24360
gtgtgctaaa	tgatatcctt	tcgagacttg	ataaagtcca	ggcggaggta	caaattgaca	24420

-continued

ggtaattac aggcagactt caaagccttc aaacctatgt aacacaacaa ctaatcaggg 24480
 ctgctgaaat cagggtctct gctaactctg ctgctactaa aatgtctgag tgtgttcttg 24540
 gacaatcaaa aagagttgac ttttgtgaa agggctacca ccttatgtcc tccccacaag 24600
 cagccccgca tgggtgtgtc ttctacatg tcacgtatgt gccatcccag gagaggaact 24660
 tcaccacagc gccagcaatt tgtcatgaag gcaaagcata cttccctcgt gaaggtgttt 24720
 ttgtgtttaa tggcacttct tggtttatta cacagaggaa cttcttttct ccacaaataa 24780
 ttactacaga caatacattt gtctcaggaa attgtgatgt cgttattggc atcattaaca 24840
 acacagttta tgatectctg caaactgagc tcgactcatt caaagaagag ctggacaagt 24900
 acttcaaaaa tcatacatca ccagatgttg atcttggcga catttcaggc attaacgctt 24960
 ctgtcgtcaa cattcaaaaa gaaattgacc gcctcaatga ggctcgtaaa aatttaaattg 25020
 aatcactcat tgaccttcaa gaattgggaa aatatgagca atatattaaa tggccttgggt 25080
 atgtttggct cggcttcatt gctggactaa ttgccatcgt catggttaca atcttgcttt 25140
 gttgcatgac tagttgttgc agttgectca aggggtcatg ctcttgtgggt tcttgcgca 25200
 agtttgatga ggatgactct gagccagttc tcaaggggtg caaattacat tacacataaa 25260
 cgaacttatg gatttgttta tgagatttt tactcttggg tcaattactg cacagccagt 25320
 aaaaattgac aatgcttctc ctgcaagtac tgttcatgct acagcaacga taccgctaca 25380
 agcctcactc cctttcggat ggcttgttat tggcgttga tttcttgctg tttttcagag 25440
 cgctacccaaa ataattgcgc tcaataaaag atggcageta gccctttata agggcttcca 25500
 gttcatttgc aatttactgc tgctatttgt taccatctat tcacatcttt tgcttgcgc 25560
 tgcaggtatg gagggcaat tttgtacct ctatgccttg atatattttc tacaatgcat 25620
 caacgcatgt agaattatta tgagatgttg gctttgttgg aagtgcaaat ccaagaacct 25680
 attactttat gatccaact actttgtttg ctggcacaca cataactatg actactgtat 25740
 accatataac agtgtcacag atacaattgt cgttactgaa ggtgacggca tttcaacacc 25800
 aaaaactcaaa gaagactacc aaattgggtg ttattctgag gatagggcact caggtgttaa 25860
 agactatgct gttgtacatg gctatttcac cgaagtttac taccagcttg agtctacaca 25920
 aattactaca gacactggta ttgaaaatgc tacattcttc atctttaaca agcttgttaa 25980
 agacccaccg aatgtgcaaa tacacacaat cgacggctct tcaggagtgg ctaatccagc 26040
 aatggatcca atttatgatg agccgacgac gactactagc gtgccttgtg aagcacaaga 26100
 aagtgagtac gaacttatgt actcattcgt ttcggaagaa acaggtagct taatagttaa 26160
 tagcgtactt ctttttcttg ctttcgtgggt attcttgcta gtcacactag ccatccttac 26220
 tgcgcttcga ttgtgtcgt actgctgcaa tattgttaac gtgagtttag taaaaccaac 26280
 ggtttacgct tactcgcgtg ttaaaaatct gaactcttct gaaggagtct ctgatcttct 26340
 ggtctaaacg aactaactat tattattatt ctgtttggaa ctttaacatt gcttatcatg 26400
 gcagacaacg gtactattac cgttgaggag cttaacaac tccctggaaca atggaaccta 26460
 gtaatagggt tcctattcct agcctggatt atgttactac aatttgccta ttctaactcg 26520
 aacaggtttt tgtacataat aaagcttgtt ttctctggc tcttggggc agtaacactt 26580
 gcttgttttg tgcttgcgct tgtctacaga attaattggg tgactggcgg gattgcgatt 26640
 gcaatggctt gtattgtagg cttgatgtgg cttagctact tcgttgcttc cttcagctg 26700
 tttgctcgtc cccgctcaat gtggctattc aaccagaaa caaacattct tctcaatgtg 26760

-continued

ctctctcggg	ggacaattgt	gaccagaccg	ctcatggaaa	gtgaacttgt	cattggtgct	26820
gtgatcattc	gtggtcactt	gcgaatggcc	ggacaccccc	tagggcgctg	tgacattaag	26880
gacctgccaa	aagagatcac	tgtggctaca	tcacgaacgc	tttcttatta	caaattagga	26940
gcgctgcagc	gtgtaggcac	tgattcaggt	tttgcctgat	acaaccgcta	ccgtattgga	27000
aactataaat	taaatacaga	ccacgccggt	agcaacgaca	atattgcttt	gctagtacag	27060
taagtgacaa	cagatgtttc	atcttgttga	cttccaggtt	acaatagcag	agatattgat	27120
tatcattatg	aggactttca	ggattgctat	ttggaatctt	gacgttataa	taagttcaat	27180
agtgagacaa	ttatttaagc	ctctaactaa	gaagaattat	toggagttag	atgatgaaga	27240
acctatggag	ttagattatc	cataaaacga	acatgaaaat	tattctcttc	ctgacattga	27300
ttgtatttac	atcttgcgag	ctatatcact	atcaggagtg	tgttagaggt	acgactgtac	27360
tactaaaaga	accttgccca	tcaggaacat	acgagggcaa	ttcaccattt	cacctcttg	27420
ctgacaataa	atttgcacta	acttgcacta	gcacacactt	tgcttttgct	tgtgctgacg	27480
gtactcgaca	tacctatcag	ctcgcgtgca	gatcagtttc	acaaaaactt	ttcatcagac	27540
aagaggaggt	tcaacaagag	ctctactcgc	cactttttct	cattgttgct	gctctagtat	27600
ttttaatact	ttgcttcacc	attaagagaa	agacagaatg	aatgagctca	ctttaattga	27660
cttctatttg	tgctttttag	cctttctgct	attccttgtt	ttaataatgc	ttattatatt	27720
ttggttttca	ctcgaaatcc	aggatctaga	agaacctgtt	accaaagtct	aaacgaacat	27780
gaaacttctc	attgttttga	cttgattttc	tctatgcagt	tgcatatgca	ctgtagtaca	27840
gcgctgtgca	tctaataaac	ctcatgtgct	tgaagatcct	tgtaaggtag	aacactaggg	27900
gtaactacta	tagcactgct	tgctttgtg	ctctaggaaa	ggttttacct	ttcatagat	27960
ggcacactat	ggttcaaaaca	tgcacaccta	atggtactat	caactgtcaa	gatccagctg	28020
gtggtgcgct	tatagctagg	tgtttgtacc	ttcatgaagg	tcaccaaact	gctgcattta	28080
gagacgtact	tgttgtttta	aataaacgaa	caaattaaaa	tgtctgataa	tggaacccaa	28140
tcaaaccaac	gtagtgcccc	cgcattaca	tttgggtggc	ccacagattc	aactgacaat	28200
aaccagaatg	gaggacgcaa	tggggcaagg	ccaaaacagc	gccgacccca	aggtttacct	28260
aataaactg	cgtcttgggt	cacagctctc	actcagcatg	gcaaggagga	acttagattc	28320
cctcgaggcc	aggcggttcc	aatcaacacc	aatagtggct	cagatgacca	aattggctac	28380
taccgaagag	ctaccgacg	agttcgtggt	ggtgacggca	aatgaaaga	gctcagcccc	28440
agatggtact	tctattacct	aggaactggc	ccagaagctt	cacttcctca	cggcgctaac	28500
aaagaaggca	tcgtatgggt	tgcaactgag	ggagccttga	atacacccaa	agaccacatt	28560
ggcacccgca	atcctaataa	caatgctgcc	accgtgctac	aacttcctca	aggacaaca	28620
ttgccaaaag	gcttctacgc	agagggaaag	agaggcggca	gtcaagcctc	ttctcgtctc	28680
tcatcacgta	gtcgcggtaa	ttcaagaaat	tcaactcctg	gcagcagtag	gggaaattct	28740
cctgctcgaa	tggctagcgg	agggtgtgaa	actgcctctg	cgctattgct	gctagacaga	28800
ttgaaccagc	ttgagagcaa	agtttctggt	aaaggccaac	aacaacaagg	ccaaactgtc	28860
actaagaaat	ctgctgctga	ggcatctaaa	aagcctcgcc	aaaaacgtac	tgccacaaaa	28920
cagtacaacg	tactcaagc	atctgggaga	cgtggtccag	aacaaaccca	aggaaatttc	28980
ggggaccaag	acctaactcag	acaaggaact	gattacaaac	attggccgca	aattgcacaa	29040
tttgcctcaa	gtgcctctgc	attctttgga	atgtcacgca	ttggcatgga	agtcacacct	29100
tcgggaacat	ggctgactta	tcattggagcc	atataattgg	atgacaaaga	tccacaattc	29160

-continued

aaagacaacg tcatactgct gaacaagcac attgacgcat acaaaacatt cccaccaaca 29220
gagcctaaaa aggacaaaaa gaaaaagact gatgaagctc agcctttgcc gcagagacaa 29280
aagaagcagc ccaactgtgac tcttcttctc gcggtgaca tggatgattt ctccagacaa 29340
cttcaaaatt ccatgagtggt agctttctgct gattcaactc aggcataaac actcatgatg 29400
accacacaag gcagatgggc tatgtaaacy ttttcgcaat tccgtttacg atacatagtc 29460
tactcttgtg cagaatgaat tctcgttaact aaacagcaca agtaggttta gttaacttta 29520
atctcacata gcaatcttta atcaatgtgt aacattaggg aggacttgaa agagccacca 29580
cattttcatc gaggccacgc ggagtacgat cgaggggtaca gtgaataatg ctaggggagag 29640
ctgcctatat ggaagagccc taatgtgtaa aattaatfff agtagtgcta tccccatgtg 29700
attttaatag cttcttagga gaatgac 29727

<210> SEQ ID NO 2
<211> LENGTH: 4382
<212> TYPE: PRT
<213> ORGANISM: Coronavirus

<400> SEQUENCE: 2

Met Glu Ser Leu Val Leu Gly Val Asn Glu Lys Thr His Val Gln Leu
1 5 10 15
Ser Leu Pro Val Leu Gln Val Arg Asp Val Leu Val Arg Gly Phe Gly
20 25 30
Asp Ser Val Glu Glu Ala Leu Ser Glu Ala Arg Glu His Leu Lys Asn
35 40 45
Gly Thr Cys Gly Leu Val Glu Leu Glu Lys Gly Val Leu Pro Gln Leu
50 55 60
Glu Gln Pro Tyr Val Phe Ile Lys Arg Ser Asp Ala Leu Ser Thr Asn
65 70 75 80
His Gly His Lys Val Val Glu Leu Val Ala Glu Met Asp Gly Ile Gln
85 90 95
Tyr Gly Arg Ser Gly Ile Thr Leu Gly Val Leu Val Pro His Val Gly
100 105 110
Glu Thr Pro Ile Ala Tyr Arg Asn Val Leu Leu Arg Lys Asn Gly Asn
115 120 125
Lys Gly Ala Gly Gly His Ser Tyr Gly Ile Asp Leu Lys Ser Tyr Asp
130 135 140
Leu Gly Asp Glu Leu Gly Thr Asp Pro Ile Glu Asp Tyr Glu Gln Asn
145 150 155 160
Trp Asn Thr Lys His Gly Ser Gly Ala Leu Arg Glu Leu Thr Arg Glu
165 170 175
Leu Asn Gly Gly Ala Val Thr Arg Tyr Val Asp Asn Asn Phe Cys Gly
180 185 190
Pro Asp Gly Tyr Pro Leu Asp Cys Ile Lys Asp Phe Leu Ala Arg Ala
195 200 205
Gly Lys Ser Met Cys Thr Leu Ser Glu Gln Leu Asp Tyr Ile Glu Ser
210 215 220
Lys Arg Gly Val Tyr Cys Cys Arg Asp His Glu His Glu Ile Ala Trp
225 230 235 240
Phe Thr Glu Arg Ser Asp Lys Ser Tyr Glu His Gln Thr Pro Phe Glu
245 250 255
Ile Lys Ser Ala Lys Lys Phe Asp Thr Phe Lys Gly Glu Cys Pro Lys
260 265 270

-continued

Phe Val Phe Pro Leu Asn Ser Lys Val Lys Val Ile Gln Pro Arg Val
 275 280 285

Glu Lys Lys Lys Thr Glu Gly Phe Met Gly Arg Ile Arg Ser Val Tyr
 290 295 300

Pro Val Ala Ser Pro Gln Glu Cys Asn Asn Met His Leu Ser Thr Leu
 305 310 315 320

Met Lys Cys Asn His Cys Asp Glu Val Ser Trp Gln Thr Cys Asp Phe
 325 330 335

Leu Lys Ala Thr Cys Glu His Cys Gly Thr Glu Asn Leu Val Ile Glu
 340 345 350

Gly Pro Thr Thr Cys Gly Tyr Leu Pro Thr Asn Ala Val Val Lys Met
 355 360 365

Pro Cys Pro Ala Cys Gln Asp Pro Glu Ile Gly Pro Glu His Ser Val
 370 375 380

Ala Asp Tyr His Asn His Ser Asn Ile Glu Thr Arg Leu Arg Lys Gly
 385 390 395 400

Gly Arg Thr Arg Cys Phe Gly Gly Cys Val Phe Ala Tyr Val Gly Cys
 405 410 415

Tyr Asn Lys Arg Ala Tyr Trp Val Pro Arg Ala Ser Ala Asp Ile Gly
 420 425 430

Ser Gly His Thr Gly Ile Thr Gly Asp Asn Val Glu Thr Leu Asn Glu
 435 440 445

Asp Leu Leu Glu Ile Leu Ser Arg Glu Arg Val Asn Ile Asn Ile Val
 450 455 460

Gly Asp Phe His Leu Asn Glu Glu Val Ala Ile Ile Leu Ala Ser Phe
 465 470 475 480

Ser Ala Ser Thr Ser Ala Phe Ile Asp Thr Ile Lys Ser Leu Asp Tyr
 485 490 495

Lys Ser Phe Lys Thr Ile Val Glu Ser Cys Gly Asn Tyr Lys Val Thr
 500 505 510

Lys Gly Lys Pro Val Lys Gly Ala Trp Asn Ile Gly Gln Gln Arg Ser
 515 520 525

Val Leu Thr Pro Leu Cys Gly Phe Pro Ser Gln Ala Ala Gly Val Ile
 530 535 540

Arg Ser Ile Phe Ala Arg Thr Leu Asp Ala Ala Asn His Ser Ile Pro
 545 550 555 560

Asp Leu Gln Arg Ala Ala Val Thr Ile Leu Asp Gly Ile Ser Glu Gln
 565 570 575

Ser Leu Arg Leu Val Asp Ala Met Val Tyr Thr Ser Asp Leu Leu Thr
 580 585 590

Asn Ser Val Ile Ile Met Ala Tyr Val Thr Gly Gly Leu Val Gln Gln
 595 600 605

Thr Ser Gln Trp Leu Ser Asn Leu Leu Gly Thr Thr Val Glu Lys Leu
 610 615 620

Arg Pro Ile Phe Glu Trp Ile Glu Ala Lys Leu Ser Ala Gly Val Glu
 625 630 635 640

Phe Leu Lys Asp Ala Trp Glu Ile Leu Lys Phe Leu Ile Thr Gly Val
 645 650 655

Phe Asp Ile Val Lys Gly Gln Ile Gln Val Ala Ser Asp Asn Ile Lys
 660 665 670

Asp Cys Val Lys Cys Phe Ile Asp Val Val Asn Lys Ala Leu Glu Met
 675 680 685

-continued

Cys Ile Asp Gln Val Thr Ile Ala Gly Ala Lys Leu Arg Ser Leu Asn
 690 695 700
 Leu Gly Glu Val Phe Ile Ala Gln Ser Lys Gly Leu Tyr Arg Gln Cys
 705 710 715 720
 Ile Arg Gly Lys Glu Gln Leu Gln Leu Leu Met Pro Leu Lys Ala Pro
 725 730 735
 Lys Glu Val Thr Phe Leu Glu Gly Asp Ser His Asp Thr Val Leu Thr
 740 745 750
 Ser Glu Glu Val Val Leu Lys Asn Gly Glu Leu Glu Ala Leu Glu Thr
 755 760 765
 Pro Val Asp Ser Phe Thr Asn Gly Ala Ile Val Gly Thr Pro Val Cys
 770 775 780
 Val Asn Gly Leu Met Leu Leu Glu Ile Lys Asp Lys Glu Gln Tyr Cys
 785 790 795 800
 Ala Leu Ser Pro Gly Leu Leu Ala Thr Asn Asn Val Phe Arg Leu Lys
 805 810 815
 Gly Gly Ala Pro Ile Lys Gly Val Thr Phe Gly Glu Asp Thr Val Trp
 820 825 830
 Glu Val Gln Gly Tyr Lys Asn Val Arg Ile Thr Phe Glu Leu Asp Glu
 835 840 845
 Arg Val Asp Lys Val Leu Asn Glu Lys Cys Ser Val Tyr Thr Val Glu
 850 855 860
 Ser Gly Thr Glu Val Thr Glu Phe Ala Cys Val Val Ala Glu Ala Val
 865 870 875 880
 Val Lys Thr Leu Gln Pro Val Ser Asp Leu Leu Thr Asn Met Gly Ile
 885 890 895
 Asp Leu Asp Glu Trp Ser Val Ala Thr Phe Tyr Leu Phe Asp Asp Ala
 900 905 910
 Gly Glu Glu Asn Phe Ser Ser Arg Met Tyr Cys Ser Phe Tyr Pro Pro
 915 920 925
 Asp Glu Glu Glu Glu Asp Asp Ala Glu Cys Glu Glu Glu Glu Ile Asp
 930 935 940
 Glu Thr Cys Glu His Glu Tyr Gly Thr Glu Asp Asp Tyr Gln Gly Leu
 945 950 955 960
 Pro Leu Glu Phe Gly Ala Ser Ala Glu Thr Val Arg Val Glu Glu Glu
 965 970 975
 Glu Glu Glu Asp Trp Leu Asp Asp Thr Thr Glu Gln Ser Glu Ile Glu
 980 985 990
 Pro Glu Pro Glu Pro Thr Pro Glu Glu Pro Val Asn Gln Phe Thr Gly
 995 1000 1005
 Tyr Leu Lys Leu Thr Asp Asn Val Ala Ile Lys Cys Val Asp Ile
 1010 1015 1020
 Val Lys Glu Ala Gln Ser Ala Asn Pro Met Val Ile Val Asn Ala
 1025 1030 1035
 Ala Asn Ile His Leu Lys His Gly Gly Gly Val Ala Gly Ala Leu
 1040 1045 1050
 Asn Lys Ala Thr Asn Gly Ala Met Gln Lys Glu Ser Asp Asp Tyr
 1055 1060 1065
 Ile Lys Leu Asn Gly Pro Leu Thr Val Gly Gly Ser Cys Leu Leu
 1070 1075 1080
 Ser Gly His Asn Leu Ala Lys Lys Cys Leu His Val Val Gly Pro
 1085 1090 1095
 Asn Leu Asn Ala Gly Glu Asp Ile Gln Leu Leu Lys Ala Ala Tyr

-continued

1100	1105	1110
Glu Asn Phe Asn Ser Gln Asp Ile Leu Leu Ala Pro Leu Leu Ser 1115 1120 1125		
Ala Gly Ile Phe Gly Ala Lys Pro Leu Gln Ser Leu Gln Val Cys 1130 1135 1140		
Val Gln Thr Val Arg Thr Gln Val Tyr Ile Ala Val Asn Asp Lys 1145 1150 1155		
Ala Leu Tyr Glu Gln Val Val Met Asp Tyr Leu Asp Asn Leu Lys 1160 1165 1170		
Pro Arg Val Glu Ala Pro Lys Gln Glu Glu Pro Pro Asn Thr Glu 1175 1180 1185		
Asp Ser Lys Thr Glu Glu Lys Ser Val Val Gln Lys Pro Val Asp 1190 1195 1200		
Val Lys Pro Lys Ile Lys Ala Cys Ile Asp Glu Val Thr Thr Thr 1205 1210 1215		
Leu Glu Glu Thr Lys Phe Leu Thr Asn Lys Leu Leu Leu Phe Ala 1220 1225 1230		
Asp Ile Asn Gly Lys Leu Tyr His Asp Ser Gln Asn Met Leu Arg 1235 1240 1245		
Gly Glu Asp Met Ser Phe Leu Glu Lys Asp Ala Pro Tyr Met Val 1250 1255 1260		
Gly Asp Val Ile Thr Ser Gly Asp Ile Thr Cys Val Val Ile Pro 1265 1270 1275		
Ser Lys Lys Ala Gly Gly Thr Thr Glu Met Leu Ser Arg Ala Leu 1280 1285 1290		
Lys Lys Val Pro Val Asp Glu Tyr Ile Thr Thr Tyr Pro Gly Gln 1295 1300 1305		
Gly Cys Ala Gly Tyr Thr Leu Glu Glu Ala Lys Thr Ala Leu Lys 1310 1315 1320		
Lys Cys Lys Ser Ala Phe Tyr Val Leu Pro Ser Glu Ala Pro Asn 1325 1330 1335		
Ala Lys Glu Glu Ile Leu Gly Thr Val Ser Trp Asn Leu Arg Glu 1340 1345 1350		
Met Leu Ala His Ala Glu Glu Thr Arg Lys Leu Met Pro Ile Cys 1355 1360 1365		
Met Asp Val Arg Ala Ile Met Ala Thr Ile Gln Arg Lys Tyr Lys 1370 1375 1380		
Gly Ile Lys Ile Gln Glu Gly Ile Val Asp Tyr Gly Val Arg Phe 1385 1390 1395		
Phe Phe Tyr Thr Ser Lys Glu Pro Val Ala Ser Ile Ile Thr Lys 1400 1405 1410		
Leu Asn Ser Leu Asn Glu Pro Leu Val Thr Met Pro Ile Gly Tyr 1415 1420 1425		
Val Thr His Gly Phe Asn Leu Glu Glu Ala Ala Arg Cys Met Arg 1430 1435 1440		
Ser Leu Lys Ala Pro Ala Val Val Ser Val Ser Ser Pro Asp Ala 1445 1450 1455		
Val Thr Thr Tyr Asn Gly Tyr Leu Thr Ser Ser Ser Lys Thr Ser 1460 1465 1470		
Glu Glu His Phe Val Glu Thr Val Ser Leu Ala Gly Ser Tyr Arg 1475 1480 1485		
Asp Trp Ser Tyr Ser Gly Gln Arg Thr Glu Leu Gly Val Glu Phe 1490 1495 1500		

-continued

Leu	Lys	Arg	Gly	Asp	Lys	Ile	Val	Tyr	His	Thr	Leu	Glu	Ser	Pro
1505						1510					1515			
Val	Glu	Phe	His	Leu	Asp	Gly	Glu	Val	Leu	Ser	Leu	Asp	Lys	Leu
1520						1525					1530			
Lys	Ser	Leu	Leu	Ser	Leu	Arg	Glu	Val	Lys	Thr	Ile	Lys	Val	Phe
1535						1540					1545			
Thr	Thr	Val	Asp	Asn	Thr	Asn	Leu	His	Thr	Gln	Leu	Val	Asp	Met
1550						1555					1560			
Ser	Met	Thr	Tyr	Gly	Gln	Gln	Phe	Gly	Pro	Thr	Tyr	Leu	Asp	Gly
1565						1570					1575			
Ala	Asp	Val	Thr	Lys	Ile	Lys	Pro	His	Val	Asn	His	Glu	Gly	Lys
1580						1585					1590			
Thr	Phe	Phe	Val	Leu	Pro	Ser	Asp	Asp	Thr	Leu	Arg	Ser	Glu	Ala
1595						1600					1605			
Phe	Glu	Tyr	Tyr	His	Thr	Leu	Asp	Glu	Ser	Phe	Leu	Gly	Arg	Tyr
1610						1615					1620			
Met	Ser	Ala	Leu	Asn	His	Thr	Lys	Lys	Trp	Lys	Phe	Pro	Gln	Val
1625						1630					1635			
Gly	Gly	Leu	Thr	Ser	Ile	Lys	Trp	Ala	Asp	Asn	Asn	Cys	Tyr	Leu
1640						1645					1650			
Ser	Ser	Val	Leu	Leu	Ala	Leu	Gln	Gln	Leu	Glu	Val	Lys	Phe	Asn
1655						1660					1665			
Ala	Pro	Ala	Leu	Gln	Glu	Ala	Tyr	Tyr	Arg	Ala	Arg	Ala	Gly	Asp
1670						1675					1680			
Ala	Ala	Asn	Phe	Cys	Ala	Leu	Ile	Leu	Ala	Tyr	Ser	Asn	Lys	Thr
1685						1690					1695			
Val	Gly	Glu	Leu	Gly	Asp	Val	Arg	Glu	Thr	Met	Thr	His	Leu	Leu
1700						1705					1710			
Gln	His	Ala	Asn	Leu	Glu	Ser	Ala	Lys	Arg	Val	Leu	Asn	Val	Val
1715						1720					1725			
Cys	Lys	His	Cys	Gly	Gln	Lys	Thr	Thr	Thr	Leu	Thr	Gly	Val	Glu
1730						1735					1740			
Ala	Val	Met	Tyr	Met	Gly	Thr	Leu	Ser	Tyr	Asp	Asn	Leu	Lys	Thr
1745						1750					1755			
Gly	Val	Ser	Ile	Pro	Cys	Val	Cys	Gly	Arg	Asp	Ala	Thr	Gln	Tyr
1760						1765					1770			
Leu	Val	Gln	Gln	Glu	Ser	Ser	Phe	Val	Met	Met	Ser	Ala	Pro	Pro
1775						1780					1785			
Ala	Glu	Tyr	Lys	Leu	Gln	Gln	Gly	Thr	Phe	Leu	Cys	Ala	Asn	Glu
1790						1795					1800			
Tyr	Thr	Gly	Asn	Tyr	Gln	Cys	Gly	His	Tyr	Thr	His	Ile	Thr	Ala
1805						1810					1815			
Lys	Glu	Thr	Leu	Tyr	Arg	Ile	Asp	Gly	Ala	His	Leu	Thr	Lys	Met
1820						1825					1830			
Ser	Glu	Tyr	Lys	Gly	Pro	Val	Thr	Asp	Val	Phe	Tyr	Lys	Glu	Thr
1835						1840					1845			
Ser	Tyr	Thr	Thr	Thr	Ile	Lys	Pro	Val	Ser	Tyr	Lys	Leu	Asp	Gly
1850						1855					1860			
Val	Thr	Tyr	Thr	Glu	Ile	Glu	Pro	Lys	Leu	Asp	Gly	Tyr	Tyr	Lys
1865						1870					1875			
Lys	Asp	Asn	Ala	Tyr	Tyr	Thr	Glu	Gln	Pro	Ile	Asp	Leu	Val	Pro
1880						1885					1890			

-continued

Thr	Gln	Pro	Leu	Pro	Asn	Ala	Ser	Phe	Asp	Asn	Phe	Lys	Leu	Thr
1895						1900					1905			
Cys	Ser	Asn	Thr	Lys	Phe	Ala	Asp	Asp	Leu	Asn	Gln	Met	Thr	Gly
1910						1915					1920			
Phe	Thr	Lys	Pro	Ala	Ser	Arg	Glu	Leu	Ser	Val	Thr	Phe	Phe	Pro
1925						1930					1935			
Asp	Leu	Asn	Gly	Asp	Val	Val	Ala	Ile	Asp	Tyr	Arg	His	Tyr	Ser
1940						1945					1950			
Ala	Ser	Phe	Lys	Lys	Gly	Ala	Lys	Leu	Leu	His	Lys	Pro	Ile	Val
1955						1960					1965			
Trp	His	Ile	Asn	Gln	Ala	Thr	Thr	Lys	Thr	Thr	Phe	Lys	Pro	Asn
1970						1975					1980			
Thr	Trp	Cys	Leu	Arg	Cys	Leu	Trp	Ser	Thr	Lys	Pro	Val	Asp	Thr
1985						1990					1995			
Ser	Asn	Ser	Phe	Glu	Val	Leu	Ala	Val	Glu	Asp	Thr	Gln	Gly	Met
2000						2005					2010			
Asp	Asn	Leu	Ala	Cys	Glu	Ser	Gln	Gln	Pro	Thr	Ser	Glu	Glu	Val
2015						2020					2025			
Val	Glu	Asn	Pro	Thr	Ile	Gln	Lys	Glu	Val	Ile	Glu	Cys	Asp	Val
2030						2035					2040			
Lys	Thr	Thr	Glu	Val	Val	Gly	Asn	Val	Ile	Leu	Lys	Pro	Ser	Asp
2045						2050					2055			
Glu	Gly	Val	Lys	Val	Thr	Gln	Glu	Leu	Gly	His	Glu	Asp	Leu	Met
2060						2065					2070			
Ala	Ala	Tyr	Val	Glu	Asn	Thr	Ser	Ile	Thr	Ile	Lys	Lys	Pro	Asn
2075						2080					2085			
Glu	Leu	Ser	Leu	Ala	Leu	Gly	Leu	Lys	Thr	Ile	Ala	Thr	His	Gly
2090						2095					2100			
Ile	Ala	Ala	Ile	Asn	Ser	Val	Pro	Trp	Ser	Lys	Ile	Leu	Ala	Tyr
2105						2110					2115			
Val	Lys	Pro	Phe	Leu	Gly	Gln	Ala	Ala	Ile	Thr	Thr	Ser	Asn	Cys
2120						2125					2130			
Ala	Lys	Arg	Leu	Ala	Gln	Arg	Val	Phe	Asn	Asn	Tyr	Met	Pro	Tyr
2135						2140					2145			
Val	Phe	Thr	Leu	Leu	Phe	Gln	Leu	Cys	Thr	Phe	Thr	Lys	Ser	Thr
2150						2155					2160			
Asn	Ser	Arg	Ile	Arg	Ala	Ser	Leu	Pro	Thr	Thr	Ile	Ala	Lys	Asn
2165						2170					2175			
Ser	Val	Lys	Ser	Val	Ala	Lys	Leu	Cys	Leu	Asp	Ala	Gly	Ile	Asn
2180						2185					2190			
Tyr	Val	Lys	Ser	Pro	Lys	Phe	Ser	Lys	Leu	Phe	Thr	Ile	Ala	Met
2195						2200					2205			
Trp	Leu	Leu	Leu	Leu	Ser	Ile	Cys	Leu	Gly	Ser	Leu	Ile	Cys	Val
2210						2215					2220			
Thr	Ala	Ala	Phe	Gly	Val	Leu	Leu	Ser	Asn	Phe	Gly	Ala	Pro	Ser
2225						2230					2235			
Tyr	Cys	Asn	Gly	Val	Arg	Glu	Leu	Tyr	Leu	Asn	Ser	Ser	Asn	Val
2240						2245					2250			
Thr	Thr	Met	Asp	Phe	Cys	Glu	Gly	Ser	Phe	Pro	Cys	Ser	Ile	Cys
2255						2260					2265			
Leu	Ser	Gly	Leu	Asp	Ser	Leu	Asp	Ser	Tyr	Pro	Ala	Leu	Glu	Thr
2270						2275					2280			
Ile	Gln	Val	Thr	Ile	Ser	Ser	Tyr	Lys	Leu	Asp	Leu	Thr	Ile	Leu

-continued

2285	2290	2295
Gly Leu Ala Ala Glu Trp Val 2300	Leu Ala Tyr Met 2305	Leu Phe Thr Lys 2310
Phe Phe Tyr Leu Leu Gly 2315	Leu Ser Ala Ile Met 2320	Gln Val Phe Phe 2325
Gly Tyr Phe Ala Ser His 2330	Phe Ile Ser Asn Ser 2335	Trp Leu Met Trp 2340
Phe Ile Ile Ser Ile Val 2345	Gln Met Ala Pro Val 2350	Ser Ala Met Val 2355
Arg Met Tyr Ile Phe Phe 2360	Ala Ser Phe Tyr Tyr 2365	Ile Trp Lys Ser 2370
Tyr Val His Ile Met Asp 2375	Gly Cys Thr Ser Ser 2380	Thr Cys Met Met 2385
Cys Tyr Lys Arg Asn Arg 2390	Ala Thr Arg Val Glu 2395	Cys Thr Thr Ile 2400
Val Asn Gly Met Lys Arg 2405	Ser Phe Tyr Val Tyr 2410	Ala Asn Gly Gly 2415
Arg Gly Phe Cys Lys Thr 2420	His Asn Trp Asn Cys 2425	Leu Asn Cys Asp 2430
Thr Phe Cys Thr Gly Ser 2435	Thr Phe Ile Ser Asp 2440	Glu Val Ala Arg 2445
Asp Leu Ser Leu Gln Phe 2450	Lys Arg Pro Ile Asn 2455	Pro Thr Asp Gln 2460
Ser Ser Tyr Ile Val Asp 2465	Ser Val Ala Val Lys 2470	Asn Gly Ala Leu 2475
His Leu Tyr Phe Asp Lys 2480	Ala Gly Gln Lys Thr 2485	Tyr Glu Arg His 2490
Pro Leu Ser His Phe Val 2495	Asn Leu Asp Asn Leu 2500	Arg Ala Asn Asn 2505
Thr Lys Gly Ser Leu Pro 2510	Ile Asn Val Ile Val 2515	Phe Asp Gly Lys 2520
Ser Lys Cys Asp Glu Ser 2525	Ala Ser Lys Ser Ala 2530	Ser Val Tyr Tyr 2535
Ser Gln Leu Met Cys Gln 2540	Pro Ile Leu Leu Leu 2545	Asp Gln Val Leu 2550
Val Ser Asp Val Gly Asp 2555	Ser Thr Glu Val Ser 2560	Val Lys Met Phe 2565
Asp Ala Tyr Val Asp Thr 2570	Phe Ser Ala Thr Phe 2575	Ser Val Pro Met 2580
Glu Lys Leu Lys Ala Leu 2585	Val Ala Thr Ala His 2590	Ser Glu Leu Ala 2595
Lys Gly Val Ala Leu Asp 2600	Gly Val Leu Ser Thr 2605	Phe Val Ser Ala 2610
Ala Arg Gln Gly Val Val 2615	Asp Thr Asp Val Asp 2620	Thr Lys Asp Val 2625
Ile Glu Cys Leu Lys Leu 2630	Ser His His Ser Asp 2635	Leu Glu Val Thr 2640
Gly Asp Ser Cys Asn Asn 2645	Phe Met Leu Thr Tyr 2650	Asn Lys Val Glu 2655
Asn Met Thr Pro Arg Asp 2660	Leu Gly Ala Cys Ile 2665	Asp Cys Asn Ala 2670
Arg His Ile Asn Ala Gln 2675	Val Ala Lys Ser His 2680	Asn Val Ser Leu 2685

-continued

Ile	Trp	Asn	Val	Lys	Asp	Tyr	Met	Ser	Leu	Ser	Glu	Gln	Leu	Arg
2690						2695					2700			
Lys	Gln	Ile	Arg	Ser	Ala	Ala	Lys	Lys	Asn	Asn	Ile	Pro	Phe	Arg
2705						2710					2715			
Leu	Thr	Cys	Ala	Thr	Thr	Arg	Gln	Val	Val	Asn	Val	Ile	Thr	Thr
2720						2725					2730			
Lys	Ile	Ser	Leu	Lys	Gly	Gly	Lys	Ile	Val	Ser	Thr	Cys	Phe	Lys
2735						2740					2745			
Leu	Met	Leu	Lys	Ala	Thr	Leu	Leu	Cys	Val	Leu	Ala	Ala	Leu	Val
2750						2755					2760			
Cys	Tyr	Ile	Val	Met	Pro	Val	His	Thr	Leu	Ser	Ile	His	Asp	Gly
2765						2770					2775			
Tyr	Thr	Asn	Glu	Ile	Ile	Gly	Tyr	Lys	Ala	Ile	Gln	Asp	Gly	Val
2780						2785					2790			
Thr	Arg	Asp	Ile	Ile	Ser	Thr	Asp	Asp	Cys	Phe	Ala	Asn	Lys	His
2795						2800					2805			
Ala	Gly	Phe	Asp	Ala	Trp	Phe	Ser	Gln	Arg	Gly	Gly	Ser	Tyr	Lys
2810						2815					2820			
Asn	Asp	Lys	Ser	Cys	Pro	Val	Val	Ala	Ala	Ile	Ile	Thr	Arg	Glu
2825						2830					2835			
Ile	Gly	Phe	Ile	Val	Pro	Gly	Leu	Pro	Gly	Thr	Val	Leu	Arg	Ala
2840						2845					2850			
Ile	Asn	Gly	Asp	Phe	Leu	His	Phe	Leu	Pro	Arg	Val	Phe	Ser	Ala
2855						2860					2865			
Val	Gly	Asn	Ile	Cys	Tyr	Thr	Pro	Ser	Lys	Leu	Ile	Glu	Tyr	Ser
2870						2875					2880			
Asp	Phe	Ala	Thr	Ser	Ala	Cys	Val	Leu	Ala	Ala	Glu	Cys	Thr	Ile
2885						2890					2895			
Phe	Lys	Asp	Ala	Met	Gly	Lys	Pro	Val	Pro	Tyr	Cys	Tyr	Asp	Thr
2900						2905					2910			
Asn	Leu	Leu	Glu	Gly	Ser	Ile	Ser	Tyr	Ser	Glu	Leu	Arg	Pro	Asp
2915						2920					2925			
Thr	Arg	Tyr	Val	Leu	Met	Asp	Gly	Ser	Ile	Ile	Gln	Phe	Pro	Asn
2930						2935					2940			
Thr	Tyr	Leu	Glu	Gly	Ser	Val	Arg	Val	Val	Thr	Thr	Phe	Asp	Ala
2945						2950					2955			
Glu	Tyr	Cys	Arg	His	Gly	Thr	Cys	Glu	Arg	Ser	Glu	Val	Gly	Ile
2960						2965					2970			
Cys	Leu	Ser	Thr	Ser	Gly	Arg	Trp	Val	Leu	Asn	Asn	Glu	His	Tyr
2975						2980					2985			
Arg	Ala	Leu	Ser	Gly	Val	Phe	Cys	Gly	Val	Asp	Ala	Met	Asn	Leu
2990						2995					3000			
Ile	Ala	Asn	Ile	Phe	Thr	Pro	Leu	Val	Gln	Pro	Val	Gly	Ala	Leu
3005						3010					3015			
Asp	Val	Ser	Ala	Ser	Val	Val	Ala	Gly	Gly	Ile	Ile	Ala	Ile	Leu
3020						3025					3030			
Val	Thr	Cys	Ala	Ala	Tyr	Tyr	Phe	Met	Lys	Phe	Arg	Arg	Val	Phe
3035						3040					3045			
Gly	Glu	Tyr	Asn	His	Val	Val	Ala	Ala	Asn	Ala	Leu	Leu	Phe	Leu
3050						3055					3060			
Met	Ser	Phe	Thr	Ile	Leu	Cys	Leu	Val	Pro	Ala	Tyr	Ser	Phe	Leu
3065						3070					3075			

-continued

Pro	Gly	Val	Tyr	Ser	Val	Phe	Tyr	Leu	Tyr	Leu	Thr	Phe	Tyr	Phe
3080						3085					3090			
Thr	Asn	Asp	Val	Ser	Phe	Leu	Ala	His	Leu	Gln	Trp	Phe	Ala	Met
3095						3100					3105			
Phe	Ser	Pro	Ile	Val	Pro	Phe	Trp	Ile	Thr	Ala	Ile	Tyr	Val	Phe
3110						3115					3120			
Cys	Ile	Ser	Leu	Lys	His	Cys	His	Trp	Phe	Phe	Asn	Asn	Tyr	Leu
3125						3130					3135			
Arg	Lys	Arg	Val	Met	Phe	Asn	Gly	Val	Thr	Phe	Ser	Thr	Phe	Glu
3140						3145					3150			
Glu	Ala	Ala	Leu	Cys	Thr	Phe	Leu	Leu	Asn	Lys	Glu	Met	Tyr	Leu
3155						3160					3165			
Lys	Leu	Arg	Ser	Glu	Thr	Leu	Leu	Pro	Leu	Thr	Gln	Tyr	Asn	Arg
3170						3175					3180			
Tyr	Leu	Ala	Leu	Tyr	Asn	Lys	Tyr	Lys	Tyr	Phe	Ser	Gly	Ala	Leu
3185						3190					3195			
Asp	Thr	Thr	Ser	Tyr	Arg	Glu	Ala	Ala	Cys	Cys	His	Leu	Ala	Lys
3200						3205					3210			
Ala	Leu	Asn	Asp	Phe	Ser	Asn	Ser	Gly	Ala	Asp	Val	Leu	Tyr	Gln
3215						3220					3225			
Pro	Pro	Gln	Thr	Ser	Ile	Thr	Ser	Ala	Val	Leu	Gln	Ser	Gly	Phe
3230						3235					3240			
Arg	Lys	Met	Ala	Phe	Pro	Ser	Gly	Lys	Val	Glu	Gly	Cys	Met	Val
3245						3250					3255			
Gln	Val	Thr	Cys	Gly	Thr	Thr	Thr	Leu	Asn	Gly	Leu	Trp	Leu	Asp
3260						3265					3270			
Asp	Thr	Val	Tyr	Cys	Pro	Arg	His	Val	Ile	Cys	Thr	Ala	Glu	Asp
3275						3280					3285			
Met	Leu	Asn	Pro	Asn	Tyr	Glu	Asp	Leu	Leu	Ile	Arg	Lys	Ser	Asn
3290						3295					3300			
His	Ser	Phe	Leu	Val	Gln	Ala	Gly	Asn	Val	Gln	Leu	Arg	Val	Ile
3305						3310					3315			
Gly	His	Ser	Met	Gln	Asn	Cys	Leu	Leu	Arg	Leu	Lys	Val	Asp	Thr
3320						3325					3330			
Ser	Asn	Pro	Lys	Thr	Pro	Lys	Tyr	Lys	Phe	Val	Arg	Ile	Gln	Pro
3335						3340					3345			
Gly	Gln	Thr	Phe	Ser	Val	Leu	Ala	Cys	Tyr	Asn	Gly	Ser	Pro	Ser
3350						3355					3360			
Gly	Val	Tyr	Gln	Cys	Ala	Met	Arg	Pro	Asn	His	Thr	Ile	Lys	Gly
3365						3370					3375			
Ser	Phe	Leu	Asn	Gly	Ser	Cys	Gly	Ser	Val	Gly	Phe	Asn	Ile	Asp
3380						3385					3390			
Tyr	Asp	Cys	Val	Ser	Phe	Cys	Tyr	Met	His	His	Met	Glu	Leu	Pro
3395						3400					3405			
Thr	Gly	Val	His	Ala	Gly	Thr	Asp	Leu	Glu	Gly	Lys	Phe	Tyr	Gly
3410						3415					3420			
Pro	Phe	Val	Asp	Arg	Gln	Thr	Ala	Gln	Ala	Ala	Gly	Thr	Asp	Thr
3425						3430					3435			
Thr	Ile	Thr	Leu	Asn	Val	Leu	Ala	Trp	Leu	Tyr	Ala	Ala	Val	Ile
3440						3445					3450			
Asn	Gly	Asp	Arg	Trp	Phe	Leu	Asn	Arg	Phe	Thr	Thr	Thr	Leu	Asn
3455						3460					3465			
Asp	Phe	Asn	Leu	Val	Ala	Met	Lys	Tyr	Asn	Tyr	Glu	Pro	Leu	Thr

-continued

Asp	Ile	Leu	Leu	Ala	Lys	Asp	Thr	Thr	Glu	Ala	Phe	Glu	Lys	Met
3875						3880					3885			
Val	Ser	Leu	Leu	Ser	Val	Leu	Leu	Ser	Met	Gln	Gly	Ala	Val	Asp
3890						3895					3900			
Ile	Asn	Arg	Leu	Cys	Glu	Glu	Met	Leu	Asp	Asn	Arg	Ala	Thr	Leu
3905						3910					3915			
Gln	Ala	Ile	Ala	Ser	Glu	Phe	Ser	Ser	Leu	Pro	Ser	Tyr	Ala	Ala
3920						3925					3930			
Tyr	Ala	Thr	Ala	Gln	Glu	Ala	Tyr	Glu	Gln	Ala	Val	Ala	Asn	Gly
3935						3940					3945			
Asp	Ser	Glu	Val	Val	Leu	Lys	Lys	Leu	Lys	Lys	Ser	Leu	Asn	Val
3950						3955					3960			
Ala	Lys	Ser	Glu	Phe	Asp	Arg	Asp	Ala	Ala	Met	Gln	Arg	Lys	Leu
3965						3970					3975			
Glu	Lys	Met	Ala	Asp	Gln	Ala	Met	Thr	Gln	Met	Tyr	Lys	Gln	Ala
3980						3985					3990			
Arg	Ser	Glu	Asp	Lys	Arg	Ala	Lys	Val	Thr	Ser	Ala	Met	Gln	Thr
3995						4000					4005			
Met	Leu	Phe	Thr	Met	Leu	Arg	Lys	Leu	Asp	Asn	Asp	Ala	Leu	Asn
4010						4015					4020			
Asn	Ile	Ile	Asn	Asn	Ala	Arg	Asp	Gly	Cys	Val	Pro	Leu	Asn	Ile
4025						4030					4035			
Ile	Pro	Leu	Thr	Thr	Ala	Ala	Lys	Leu	Met	Val	Val	Val	Pro	Asp
4040						4045					4050			
Tyr	Gly	Thr	Tyr	Lys	Asn	Thr	Cys	Asp	Gly	Asn	Thr	Phe	Thr	Tyr
4055						4060					4065			
Ala	Ser	Ala	Leu	Trp	Glu	Ile	Gln	Gln	Val	Val	Asp	Ala	Asp	Ser
4070						4075					4080			
Lys	Ile	Val	Gln	Leu	Ser	Glu	Ile	Asn	Met	Asp	Asn	Ser	Pro	Asn
4085						4090					4095			
Leu	Ala	Trp	Pro	Leu	Ile	Val	Thr	Ala	Leu	Arg	Ala	Asn	Ser	Ala
4100						4105					4110			
Val	Lys	Leu	Gln	Asn	Asn	Glu	Leu	Ser	Pro	Val	Ala	Leu	Arg	Gln
4115						4120					4125			
Met	Ser	Cys	Ala	Ala	Gly	Thr	Thr	Gln	Thr	Ala	Cys	Thr	Asp	Asp
4130						4135					4140			
Asn	Ala	Leu	Ala	Tyr	Tyr	Asn	Asn	Ser	Lys	Gly	Gly	Arg	Phe	Val
4145						4150					4155			
Leu	Ala	Leu	Leu	Ser	Asp	His	Gln	Asp	Leu	Lys	Trp	Ala	Arg	Phe
4160						4165					4170			
Pro	Lys	Ser	Asp	Gly	Thr	Gly	Thr	Ile	Tyr	Thr	Glu	Leu	Glu	Pro
4175						4180					4185			
Pro	Cys	Arg	Phe	Val	Thr	Asp	Thr	Pro	Lys	Gly	Pro	Lys	Val	Lys
4190						4195					4200			
Tyr	Leu	Tyr	Phe	Ile	Lys	Gly	Leu	Asn	Asn	Leu	Asn	Arg	Gly	Met
4205						4210					4215			
Val	Leu	Gly	Ser	Leu	Ala	Ala	Thr	Val	Arg	Leu	Gln	Ala	Gly	Asn
4220						4225					4230			
Ala	Thr	Glu	Val	Pro	Ala	Asn	Ser	Thr	Val	Leu	Ser	Phe	Cys	Ala
4235						4240					4245			
Phe	Ala	Val	Asp	Pro	Ala	Lys	Ala	Tyr	Lys	Asp	Tyr	Leu	Ala	Ser
4250						4255					4260			

-continued

Gly Gly Gln Pro Ile Thr Asn Cys Val Lys Met Leu Cys Thr His
4265 4270 4275

Thr Gly Thr Gly Gln Ala Ile Thr Val Thr Pro Glu Ala Asn Met
4280 4285 4290

Asp Gln Glu Ser Phe Gly Gly Ala Ser Cys Cys Leu Tyr Cys Arg
4295 4300 4305

Cys His Ile Asp His Pro Asn Pro Lys Gly Phe Cys Asp Leu Lys
4310 4315 4320

Gly Lys Tyr Val Gln Ile Pro Thr Thr Cys Ala Asn Asp Pro Val
4325 4330 4335

Gly Phe Thr Leu Arg Asn Thr Val Cys Thr Val Cys Gly Met Trp
4340 4345 4350

Lys Gly Tyr Gly Cys Ser Cys Asp Gln Leu Arg Glu Pro Leu Met
4355 4360 4365

Gln Ser Ala Asp Ala Ser Thr Phe Leu Asn Gly Phe Ala Val
4370 4375 4380

<210> SEQ ID NO 3

<211> LENGTH: 2695

<212> TYPE: PRT

<213> ORGANISM: Coronavirus

<400> SEQUENCE: 3

Arg Val Cys Gly Val Ser Ala Ala Arg Leu Thr Pro Cys Gly Thr Gly
1 5 10 15

Thr Ser Thr Asp Val Val Tyr Arg Ala Phe Asp Ile Tyr Asn Glu Lys
20 25 30

Val Ala Gly Phe Ala Lys Phe Leu Lys Thr Asn Cys Cys Arg Phe Gln
35 40 45

Glu Lys Asp Glu Glu Gly Asn Leu Leu Asp Ser Tyr Phe Val Val Lys
50 55 60

Arg His Thr Met Ser Asn Tyr Gln His Glu Glu Thr Ile Tyr Asn Leu
65 70 75 80

Val Lys Asp Cys Pro Ala Val Ala Val His Asp Phe Phe Lys Phe Arg
85 90 95

Val Asp Gly Asp Met Val Pro His Ile Ser Arg Gln Arg Leu Thr Lys
100 105 110

Tyr Thr Met Ala Asp Leu Val Tyr Ala Leu Arg His Phe Asp Glu Gly
115 120 125

Asn Cys Asp Thr Leu Lys Glu Ile Leu Val Thr Tyr Asn Cys Cys Asp
130 135 140

Asp Asp Tyr Phe Asn Lys Lys Asp Trp Tyr Asp Phe Val Glu Asn Pro
145 150 155 160

Asp Ile Leu Arg Val Tyr Ala Asn Leu Gly Glu Arg Val Arg Gln Ser
165 170 175

Leu Leu Lys Thr Val Gln Phe Cys Asp Ala Met Arg Asp Ala Gly Ile
180 185 190

Val Gly Val Leu Thr Leu Asp Asn Gln Asp Leu Asn Gly Asn Trp Tyr
195 200 205

Asp Phe Gly Asp Phe Val Gln Val Ala Pro Gly Cys Gly Val Pro Ile
210 215 220

Val Asp Ser Tyr Tyr Ser Leu Leu Met Pro Ile Leu Thr Leu Thr Arg
225 230 235 240

Ala Leu Ala Ala Glu Ser His Met Asp Ala Asp Leu Ala Lys Pro Leu
245 250 255

-continued

Ile Lys Trp Asp Leu Leu Lys Tyr Asp Phe Thr Glu Glu Arg Leu Cys
 260 265 270

Leu Phe Asp Arg Tyr Phe Lys Tyr Trp Asp Gln Thr Tyr His Pro Asn
 275 280 285

Cys Ile Asn Cys Leu Asp Asp Arg Cys Ile Leu His Cys Ala Asn Phe
 290 295 300

Asn Val Leu Phe Ser Thr Val Phe Pro Pro Thr Ser Phe Gly Pro Leu
 305 310 315 320

Val Arg Lys Ile Phe Val Asp Gly Val Pro Phe Val Val Ser Thr Gly
 325 330 335

Tyr His Phe Arg Glu Leu Gly Val Val His Asn Gln Asp Val Asn Leu
 340 345 350

His Ser Ser Arg Leu Ser Phe Lys Glu Leu Leu Val Tyr Ala Ala Asp
 355 360 365

Pro Ala Met His Ala Ala Ser Gly Asn Leu Leu Leu Asp Lys Arg Thr
 370 375 380

Thr Cys Phe Ser Val Ala Ala Leu Thr Asn Asn Val Ala Phe Gln Thr
 385 390 395 400

Val Lys Pro Gly Asn Phe Asn Lys Asp Phe Tyr Asp Phe Ala Val Ser
 405 410 415

Lys Gly Phe Phe Lys Glu Gly Ser Ser Val Glu Leu Lys His Phe Phe
 420 425 430

Phe Ala Gln Asp Gly Asn Ala Ala Ile Ser Asp Tyr Asp Tyr Tyr Arg
 435 440 445

Tyr Asn Leu Pro Thr Met Cys Asp Ile Arg Gln Leu Leu Phe Val Val
 450 455 460

Glu Val Val Asp Lys Tyr Phe Asp Cys Tyr Asp Gly Gly Cys Ile Asn
 465 470 475 480

Ala Asn Gln Val Ile Val Asn Asn Leu Asp Lys Ser Ala Gly Phe Pro
 485 490 495

Phe Asn Lys Trp Gly Lys Ala Arg Leu Tyr Tyr Asp Ser Met Ser Tyr
 500 505 510

Glu Asp Gln Asp Ala Leu Phe Ala Tyr Thr Lys Arg Asn Val Ile Pro
 515 520 525

Thr Ile Thr Gln Met Asn Leu Lys Tyr Ala Ile Ser Ala Lys Asn Arg
 530 535 540

Ala Arg Thr Val Ala Gly Val Ser Ile Cys Ser Thr Met Thr Asn Arg
 545 550 555 560

Gln Phe His Gln Lys Leu Leu Lys Ser Ile Ala Ala Thr Arg Gly Ala
 565 570 575

Thr Val Val Ile Gly Thr Ser Lys Phe Tyr Gly Gly Trp His Asn Met
 580 585 590

Leu Lys Thr Val Tyr Ser Asp Val Glu Thr Pro His Leu Met Gly Trp
 595 600 605

Asp Tyr Pro Lys Cys Asp Arg Ala Met Pro Asn Met Leu Arg Ile Met
 610 615 620

Ala Ser Leu Val Leu Ala Arg Lys His Asn Thr Cys Cys Asn Leu Ser
 625 630 635 640

His Arg Phe Tyr Arg Leu Ala Asn Glu Cys Ala Gln Val Leu Ser Glu
 645 650 655

Met Val Met Cys Gly Gly Ser Leu Tyr Val Lys Pro Gly Gly Thr Ser
 660 665 670

-continued

Ser Gly Asp Ala Thr Thr Ala Tyr Ala Asn Ser Val Phe Asn Ile Cys
 675 680 685

Gln Ala Val Thr Ala Asn Val Asn Ala Leu Leu Ser Thr Asp Gly Asn
 690 695 700

Lys Ile Ala Asp Lys Tyr Val Arg Asn Leu Gln His Arg Leu Tyr Glu
 705 710 715 720

Cys Leu Tyr Arg Asn Arg Asp Val Asp His Glu Phe Val Asp Glu Phe
 725 730 735

Tyr Ala Tyr Leu Arg Lys His Phe Ser Met Met Ile Leu Ser Asp Asp
 740 745 750

Ala Val Val Cys Tyr Asn Ser Asn Tyr Ala Ala Gln Gly Leu Val Ala
 755 760 765

Ser Ile Lys Asn Phe Lys Ala Val Leu Tyr Tyr Gln Asn Asn Val Phe
 770 775 780

Met Ser Glu Ala Lys Cys Trp Thr Glu Thr Asp Leu Thr Lys Gly Pro
 785 790 795 800

His Glu Phe Cys Ser Gln His Thr Met Leu Val Lys Gln Gly Asp Asp
 805 810 815

Tyr Val Tyr Leu Pro Tyr Pro Asp Pro Ser Arg Ile Leu Gly Ala Gly
 820 825 830

Cys Phe Val Asp Asp Ile Val Lys Thr Asp Gly Thr Leu Met Ile Glu
 835 840 845

Arg Phe Val Ser Leu Ala Ile Asp Ala Tyr Pro Leu Thr Lys His Pro
 850 855 860

Asn Gln Glu Tyr Ala Asp Val Phe His Leu Tyr Leu Gln Tyr Ile Arg
 865 870 875 880

Lys Leu His Asp Glu Leu Thr Gly His Met Leu Asp Met Tyr Ser Val
 885 890 895

Met Leu Thr Asn Asp Asn Thr Ser Arg Tyr Trp Glu Pro Glu Phe Tyr
 900 905 910

Glu Ala Met Tyr Thr Pro His Thr Val Leu Gln Ala Val Gly Ala Cys
 915 920 925

Val Leu Cys Asn Ser Gln Thr Ser Leu Arg Cys Gly Ala Cys Ile Arg
 930 935 940

Arg Pro Phe Leu Cys Cys Lys Cys Cys Tyr Asp His Val Ile Ser Thr
 945 950 955 960

Ser His Lys Leu Val Leu Ser Val Asn Pro Tyr Val Cys Asn Ala Pro
 965 970 975

Gly Cys Asp Val Thr Asp Val Thr Gln Leu Tyr Leu Gly Gly Met Ser
 980 985 990

Tyr Tyr Cys Lys Ser His Lys Pro Pro Ile Ser Phe Pro Leu Cys Ala
 995 1000 1005

Asn Gly Gln Val Phe Gly Leu Tyr Lys Asn Thr Cys Val Gly Ser
 1010 1015 1020

Asp Asn Val Thr Asp Phe Asn Ala Ile Ala Thr Cys Asp Trp Thr
 1025 1030 1035

Asn Ala Gly Asp Tyr Ile Leu Ala Asn Thr Cys Thr Glu Arg Leu
 1040 1045 1050

Lys Leu Phe Ala Ala Glu Thr Leu Lys Ala Thr Glu Glu Thr Phe
 1055 1060 1065

Lys Leu Ser Tyr Gly Ile Ala Thr Val Arg Glu Val Leu Ser Asp
 1070 1075 1080

Arg Glu Leu His Leu Ser Trp Glu Val Gly Lys Pro Arg Pro Pro

-continued

1085		1090		1095
Leu Asn Arg Asn Tyr Val Phe Thr Gly Tyr Arg Val Thr Lys Asn 1100		1105		1110
Ser Lys Val Gln Ile Gly Glu Tyr Thr Phe Glu Lys Gly Asp Tyr 1115		1120		1125
Gly Asp Ala Val Val Tyr Arg Gly Thr Thr Thr Tyr Lys Leu Asn 1130		1135		1140
Val Gly Asp Tyr Phe Val Leu Thr Ser His Thr Val Met Pro Leu 1145		1150		1155
Ser Ala Pro Thr Leu Val Pro Gln Glu His Tyr Val Arg Ile Thr 1160		1165		1170
Gly Leu Tyr Pro Thr Leu Asn Ile Ser Asp Glu Phe Ser Ser Asn 1175		1180		1185
Val Ala Asn Tyr Gln Lys Val Gly Met Gln Lys Tyr Ser Thr Leu 1190		1195		1200
Gln Gly Pro Pro Gly Thr Gly Lys Ser His Phe Ala Ile Gly Leu 1205		1210		1215
Ala Leu Tyr Tyr Pro Ser Ala Arg Ile Val Tyr Thr Ala Cys Ser 1220		1225		1230
His Ala Ala Val Asp Ala Leu Cys Glu Lys Ala Leu Lys Tyr Leu 1235		1240		1245
Pro Ile Asp Lys Cys Ser Arg Ile Ile Pro Ala Arg Ala Arg Val 1250		1255		1260
Glu Cys Phe Asp Lys Phe Lys Val Asn Ser Thr Leu Glu Gln Tyr 1265		1270		1275
Val Phe Cys Thr Val Asn Ala Leu Pro Glu Thr Thr Ala Asp Ile 1280		1285		1290
Val Val Phe Asp Glu Ile Ser Met Ala Thr Asn Tyr Asp Leu Ser 1295		1300		1305
Val Val Asn Ala Arg Leu Arg Ala Lys His Tyr Val Tyr Ile Gly 1310		1315		1320
Asp Pro Ala Gln Leu Pro Ala Pro Arg Thr Leu Leu Thr Lys Gly 1325		1330		1335
Thr Leu Glu Pro Glu Tyr Phe Asn Ser Val Cys Arg Leu Met Lys 1340		1345		1350
Thr Ile Gly Pro Asp Met Phe Leu Gly Thr Cys Arg Arg Cys Pro 1355		1360		1365
Ala Glu Ile Val Asp Thr Val Ser Ala Leu Val Tyr Asp Asn Lys 1370		1375		1380
Leu Lys Ala His Lys Asp Lys Ser Ala Gln Cys Phe Lys Met Phe 1385		1390		1395
Tyr Lys Gly Val Ile Thr His Asp Val Ser Ser Ala Ile Asn Arg 1400		1405		1410
Pro Gln Ile Gly Val Val Arg Glu Phe Leu Thr Arg Asn Pro Ala 1415		1420		1425
Trp Arg Lys Ala Val Phe Ile Ser Pro Tyr Asn Ser Gln Asn Ala 1430		1435		1440
Val Ala Ser Lys Ile Leu Gly Leu Pro Thr Gln Thr Val Asp Ser 1445		1450		1455
Ser Gln Gly Ser Glu Tyr Asp Tyr Val Ile Phe Thr Gln Thr Thr 1460		1465		1470
Glu Thr Ala His Ser Cys Asn Val Asn Arg Phe Asn Val Ala Ile 1475		1480		1485

-continued

Thr	Arg	Ala	Lys	Ile	Gly	Ile	Leu	Cys	Ile	Met	Ser	Asp	Arg	Asp
1490						1495						1500		
Leu	Tyr	Asp	Lys	Leu	Gln	Phe	Thr	Ser	Leu	Glu	Ile	Pro	Arg	Arg
1505						1510						1515		
Asn	Val	Ala	Thr	Leu	Gln	Ala	Glu	Asn	Val	Thr	Gly	Leu	Phe	Lys
1520						1525						1530		
Asp	Cys	Ser	Lys	Ile	Ile	Thr	Gly	Leu	His	Pro	Thr	Gln	Ala	Pro
1535						1540						1545		
Thr	His	Leu	Ser	Val	Asp	Ile	Lys	Phe	Lys	Thr	Glu	Gly	Leu	Cys
1550						1555						1560		
Val	Asp	Ile	Pro	Gly	Ile	Pro	Lys	Asp	Met	Thr	Tyr	Arg	Arg	Leu
1565						1570						1575		
Ile	Ser	Met	Met	Gly	Phe	Lys	Met	Asn	Tyr	Gln	Val	Asn	Gly	Tyr
1580						1585						1590		
Pro	Asn	Met	Phe	Ile	Thr	Arg	Glu	Glu	Ala	Ile	Arg	His	Val	Arg
1595						1600						1605		
Ala	Trp	Ile	Gly	Phe	Asp	Val	Glu	Gly	Cys	His	Ala	Thr	Arg	Asp
1610						1615						1620		
Ala	Val	Gly	Thr	Asn	Leu	Pro	Leu	Gln	Leu	Gly	Phe	Ser	Thr	Gly
1625						1630						1635		
Val	Asn	Leu	Val	Ala	Val	Pro	Thr	Gly	Tyr	Val	Asp	Thr	Glu	Asn
1640						1645						1650		
Asn	Thr	Glu	Phe	Thr	Arg	Val	Asn	Ala	Lys	Pro	Pro	Pro	Gly	Asp
1655						1660						1665		
Gln	Phe	Lys	His	Leu	Ile	Pro	Leu	Met	Tyr	Lys	Gly	Leu	Pro	Trp
1670						1675						1680		
Asn	Val	Val	Arg	Ile	Lys	Ile	Val	Gln	Met	Leu	Ser	Asp	Thr	Leu
1685						1690						1695		
Lys	Gly	Leu	Ser	Asp	Arg	Val	Val	Phe	Val	Leu	Trp	Ala	His	Gly
1700						1705						1710		
Phe	Glu	Leu	Thr	Ser	Met	Lys	Tyr	Phe	Val	Lys	Ile	Gly	Pro	Glu
1715						1720						1725		
Arg	Thr	Cys	Cys	Leu	Cys	Asp	Lys	Arg	Ala	Thr	Cys	Phe	Ser	Thr
1730						1735						1740		
Ser	Ser	Asp	Thr	Tyr	Ala	Cys	Trp	Asn	His	Ser	Val	Gly	Phe	Asp
1745						1750						1755		
Tyr	Val	Tyr	Asn	Pro	Phe	Met	Ile	Asp	Val	Gln	Gln	Trp	Gly	Phe
1760						1765						1770		
Thr	Gly	Asn	Leu	Gln	Ser	Asn	His	Asp	Gln	His	Cys	Gln	Val	His
1775						1780						1785		
Gly	Asn	Ala	His	Val	Ala	Ser	Cys	Asp	Ala	Ile	Met	Thr	Arg	Cys
1790						1795						1800		
Leu	Ala	Val	His	Glu	Cys	Phe	Val	Lys	Arg	Val	Asp	Trp	Ser	Val
1805						1810						1815		
Glu	Tyr	Pro	Ile	Ile	Gly	Asp	Glu	Leu	Arg	Val	Asn	Ser	Ala	Cys
1820						1825						1830		
Arg	Lys	Val	Gln	His	Met	Val	Val	Lys	Ser	Ala	Leu	Leu	Ala	Asp
1835						1840						1845		
Lys	Phe	Pro	Val	Leu	His	Asp	Ile	Gly	Asn	Pro	Lys	Ala	Ile	Lys
1850						1855						1860		
Cys	Val	Pro	Gln	Ala	Glu	Val	Glu	Trp	Lys	Phe	Tyr	Asp	Ala	Gln
1865						1870						1875		

-continued

Pro	Cys	Ser	Asp	Lys	Ala	Tyr	Lys	Ile	Glu	Glu	Leu	Phe	Tyr	Ser
	1880					1885					1890			
Tyr	Ala	Thr	His	His	Asp	Lys	Phe	Thr	Asp	Gly	Val	Cys	Leu	Phe
	1895					1900					1905			
Trp	Asn	Cys	Asn	Val	Asp	Arg	Tyr	Pro	Ala	Asn	Ala	Ile	Val	Cys
	1910					1915					1920			
Arg	Phe	Asp	Thr	Arg	Val	Leu	Ser	Asn	Leu	Asn	Leu	Pro	Gly	Cys
	1925					1930					1935			
Asp	Gly	Gly	Ser	Leu	Tyr	Val	Asn	Lys	His	Ala	Phe	His	Thr	Pro
	1940					1945					1950			
Ala	Phe	Asp	Lys	Ser	Ala	Phe	Thr	Asn	Leu	Lys	Gln	Leu	Pro	Phe
	1955					1960					1965			
Phe	Tyr	Tyr	Ser	Asp	Ser	Pro	Cys	Glu	Ser	His	Gly	Lys	Gln	Val
	1970					1975					1980			
Val	Ser	Asp	Ile	Asp	Tyr	Val	Pro	Leu	Lys	Ser	Ala	Thr	Cys	Ile
	1985					1990					1995			
Thr	Arg	Cys	Asn	Leu	Gly	Gly	Ala	Val	Cys	Arg	His	His	Ala	Asn
	2000					2005					2010			
Glu	Tyr	Arg	Gln	Tyr	Leu	Asp	Ala	Tyr	Asn	Met	Met	Ile	Ser	Ala
	2015					2020					2025			
Gly	Phe	Ser	Leu	Trp	Ile	Tyr	Lys	Gln	Phe	Asp	Thr	Tyr	Asn	Leu
	2030					2035					2040			
Trp	Asn	Thr	Phe	Thr	Arg	Leu	Gln	Ser	Leu	Glu	Asn	Val	Ala	Tyr
	2045					2050					2055			
Asn	Val	Val	Asn	Lys	Gly	His	Phe	Asp	Gly	His	Ala	Gly	Glu	Ala
	2060					2065					2070			
Pro	Val	Ser	Ile	Ile	Asn	Asn	Ala	Val	Tyr	Thr	Lys	Val	Asp	Gly
	2075					2080					2085			
Ile	Asp	Val	Glu	Ile	Phe	Glu	Asn	Lys	Thr	Thr	Leu	Pro	Val	Asn
	2090					2095					2100			
Val	Ala	Phe	Glu	Leu	Trp	Ala	Lys	Arg	Asn	Ile	Lys	Pro	Val	Pro
	2105					2110					2115			
Glu	Ile	Lys	Ile	Leu	Asn	Asn	Leu	Gly	Val	Asp	Ile	Ala	Ala	Asn
	2120					2125					2130			
Thr	Val	Ile	Trp	Asp	Tyr	Lys	Arg	Glu	Ala	Pro	Ala	His	Val	Ser
	2135					2140					2145			
Thr	Ile	Gly	Val	Cys	Thr	Met	Thr	Asp	Ile	Ala	Lys	Lys	Pro	Thr
	2150					2155					2160			
Glu	Ser	Ala	Cys	Ser	Ser	Leu	Thr	Val	Leu	Phe	Asp	Gly	Arg	Val
	2165					2170					2175			
Glu	Gly	Gln	Val	Asp	Leu	Phe	Arg	Asn	Ala	Arg	Asn	Gly	Val	Leu
	2180					2185					2190			
Ile	Thr	Glu	Gly	Ser	Val	Lys	Gly	Leu	Thr	Pro	Ser	Lys	Gly	Pro
	2195					2200					2205			
Ala	Gln	Ala	Ser	Val	Asn	Gly	Val	Thr	Leu	Ile	Gly	Glu	Ser	Val
	2210					2215					2220			
Lys	Thr	Gln	Phe	Asn	Tyr	Phe	Lys	Lys	Val	Asp	Gly	Ile	Ile	Gln
	2225					2230					2235			
Gln	Leu	Pro	Glu	Thr	Tyr	Phe	Thr	Gln	Ser	Arg	Asp	Leu	Glu	Asp
	2240					2245					2250			
Phe	Lys	Pro	Arg	Ser	Gln	Met	Glu	Thr	Asp	Phe	Leu	Glu	Leu	Ala
	2255					2260					2265			
Met	Asp	Glu	Phe	Ile	Gln	Arg	Tyr	Lys	Leu	Glu	Gly	Tyr	Ala	Phe

-continued

2270	2275	2280
Glu His Ile Val Tyr Gly Asp Phe Ser His Gly Gln Leu Gly Gly 2285 2290 2295		
Leu His Leu Met Ile Gly Leu Ala Lys Arg Ser Gln Asp Ser Pro 2300 2305 2310		
Leu Lys Leu Glu Asp Phe Ile Pro Met Asp Ser Thr Val Lys Asn 2315 2320 2325		
Tyr Phe Ile Thr Asp Ala Gln Thr Gly Ser Ser Lys Cys Val Cys 2330 2335 2340		
Ser Val Ile Asp Leu Leu Leu Asp Asp Phe Val Glu Ile Ile Lys 2345 2350 2355		
Ser Gln Asp Leu Ser Val Ile Ser Lys Val Val Lys Val Thr Ile 2360 2365 2370		
Asp Tyr Ala Glu Ile Ser Phe Met Leu Trp Cys Lys Asp Gly His 2375 2380 2385		
Val Glu Thr Phe Tyr Pro Lys Leu Gln Ala Ser Gln Ala Trp Gln 2390 2395 2400		
Pro Gly Val Ala Met Pro Asn Leu Tyr Lys Met Gln Arg Met Leu 2405 2410 2415		
Leu Glu Lys Cys Asp Leu Gln Asn Tyr Gly Glu Asn Ala Val Ile 2420 2425 2430		
Pro Lys Gly Ile Met Met Asn Val Ala Lys Tyr Thr Gln Leu Cys 2435 2440 2445		
Gln Tyr Leu Asn Thr Leu Thr Leu Ala Val Pro Tyr Asn Met Arg 2450 2455 2460		
Val Ile His Phe Gly Ala Gly Ser Asp Lys Gly Val Ala Pro Gly 2465 2470 2475		
Thr Ala Val Leu Arg Gln Trp Leu Pro Thr Gly Thr Leu Leu Val 2480 2485 2490		
Asp Ser Asp Leu Asn Asp Phe Val Ser Asp Ala Asp Ser Thr Leu 2495 2500 2505		
Ile Gly Asp Cys Ala Thr Val His Thr Ala Asn Lys Trp Asp Leu 2510 2515 2520		
Ile Ile Ser Asp Met Tyr Asp Pro Arg Thr Lys His Val Thr Lys 2525 2530 2535		
Glu Asn Asp Ser Lys Glu Gly Phe Phe Thr Tyr Leu Cys Gly Phe 2540 2545 2550		
Ile Lys Gln Lys Leu Ala Leu Gly Gly Ser Ile Ala Val Lys Ile 2555 2560 2565		
Thr Glu His Ser Trp Asn Ala Asp Leu Tyr Lys Leu Met Gly His 2570 2575 2580		
Phe Ser Trp Trp Thr Ala Phe Val Thr Asn Val Asn Ala Ser Ser 2585 2590 2595		
Ser Glu Ala Phe Leu Ile Gly Ala Asn Tyr Leu Gly Lys Pro Lys 2600 2605 2610		
Glu Gln Ile Asp Gly Tyr Thr Met His Ala Asn Tyr Ile Phe Trp 2615 2620 2625		
Arg Asn Thr Asn Pro Ile Gln Leu Ser Ser Tyr Ser Leu Phe Asp 2630 2635 2640		
Met Ser Lys Phe Pro Leu Lys Leu Arg Gly Thr Ala Val Met Ser 2645 2650 2655		
Leu Lys Glu Asn Gln Ile Asn Asp Met Ile Tyr Ser Leu Leu Glu 2660 2665 2670		

-continued

Lys Gly Arg Leu Ile Ile Arg Glu Asn Asn Arg Val Val Val Ser
 2675 2680 2685

Ser Asp Ile Leu Val Asn Asn
 2690 2695

<210> SEQ ID NO 4
 <211> LENGTH: 1255
 <212> TYPE: PRT
 <213> ORGANISM: Coronavirus

<400> SEQUENCE: 4

Met Phe Ile Phe Leu Leu Phe Leu Thr Leu Thr Ser Gly Ser Asp Leu
 1 5 10 15

Asp Arg Cys Thr Thr Phe Asp Asp Val Gln Ala Pro Asn Tyr Thr Gln
 20 25 30

His Thr Ser Ser Met Arg Gly Val Tyr Tyr Pro Asp Glu Ile Phe Arg
 35 40 45

Ser Asp Thr Leu Tyr Leu Thr Gln Asp Leu Phe Leu Pro Phe Tyr Ser
 50 55 60

Asn Val Thr Gly Phe His Thr Ile Asn His Thr Phe Gly Asn Pro Val
 65 70 75 80

Ile Pro Phe Lys Asp Gly Ile Tyr Phe Ala Ala Thr Glu Lys Ser Asn
 85 90 95

Val Val Arg Gly Trp Val Phe Gly Ser Thr Met Asn Asn Lys Ser Gln
 100 105 110

Ser Val Ile Ile Ile Asn Asn Ser Thr Asn Val Val Ile Arg Ala Cys
 115 120 125

Asn Phe Glu Leu Cys Asp Asn Pro Phe Phe Ala Val Ser Lys Pro Met
 130 135 140

Gly Thr Gln Thr His Thr Met Ile Phe Asp Asn Ala Phe Asn Cys Thr
 145 150 155 160

Phe Glu Tyr Ile Ser Asp Ala Phe Ser Leu Asp Val Ser Glu Lys Ser
 165 170 175

Gly Asn Phe Lys His Leu Arg Glu Phe Val Phe Lys Asn Lys Asp Gly
 180 185 190

Phe Leu Tyr Val Tyr Lys Gly Tyr Gln Pro Ile Asp Val Val Arg Asp
 195 200 205

Leu Pro Ser Gly Phe Asn Thr Leu Lys Pro Ile Phe Lys Leu Pro Leu
 210 215 220

Gly Ile Asn Ile Thr Asn Phe Arg Ala Ile Leu Thr Ala Phe Ser Pro
 225 230 235 240

Ala Gln Asp Ile Trp Gly Thr Ser Ala Ala Ala Tyr Phe Val Gly Tyr
 245 250 255

Leu Lys Pro Thr Thr Phe Met Leu Lys Tyr Asp Glu Asn Gly Thr Ile
 260 265 270

Thr Asp Ala Val Asp Cys Ser Gln Asn Pro Leu Ala Glu Leu Lys Cys
 275 280 285

Ser Val Lys Ser Phe Glu Ile Asp Lys Gly Ile Tyr Gln Thr Ser Asn
 290 295 300

Phe Arg Val Val Pro Ser Gly Asp Val Val Arg Phe Pro Asn Ile Thr
 305 310 315 320

Asn Leu Cys Pro Phe Gly Glu Val Phe Asn Ala Thr Lys Phe Pro Ser
 325 330 335

Val Tyr Ala Trp Glu Arg Lys Lys Ile Ser Asn Cys Val Ala Asp Tyr

-continued

340					345					350					
Ser	Val	Leu	Tyr	Asn	Ser	Thr	Phe	Phe	Ser	Thr	Phe	Lys	Cys	Tyr	Gly
		355					360					365			
Val	Ser	Ala	Thr	Lys	Leu	Asn	Asp	Leu	Cys	Phe	Ser	Asn	Val	Tyr	Ala
		370				375					380				
Asp	Ser	Phe	Val	Val	Lys	Gly	Asp	Asp	Val	Arg	Gln	Ile	Ala	Pro	Gly
		385			390					395					400
Gln	Thr	Gly	Val	Ile	Ala	Asp	Tyr	Asn	Tyr	Lys	Leu	Pro	Asp	Asp	Phe
				405					410					415	
Met	Gly	Cys	Val	Leu	Ala	Trp	Asn	Thr	Arg	Asn	Ile	Asp	Ala	Thr	Ser
			420					425					430		
Thr	Gly	Asn	Tyr	Asn	Tyr	Lys	Tyr	Arg	Tyr	Leu	Arg	His	Gly	Lys	Leu
		435					440					445			
Arg	Pro	Phe	Glu	Arg	Asp	Ile	Ser	Asn	Val	Pro	Phe	Ser	Pro	Asp	Gly
		450				455					460				
Lys	Pro	Cys	Thr	Pro	Pro	Ala	Leu	Asn	Cys	Tyr	Trp	Pro	Leu	Asn	Asp
		465			470					475					480
Tyr	Gly	Phe	Tyr	Thr	Thr	Thr	Gly	Ile	Gly	Tyr	Gln	Pro	Tyr	Arg	Val
				485					490					495	
Val	Val	Leu	Ser	Phe	Glu	Leu	Leu	Asn	Ala	Pro	Ala	Thr	Val	Cys	Gly
				500				505					510		
Pro	Lys	Leu	Ser	Thr	Asp	Leu	Ile	Lys	Asn	Gln	Cys	Val	Asn	Phe	Asn
		515					520					525			
Phe	Asn	Gly	Leu	Thr	Gly	Thr	Gly	Val	Leu	Thr	Pro	Ser	Ser	Lys	Arg
		530				535					540				
Phe	Gln	Pro	Phe	Gln	Gln	Phe	Gly	Arg	Asp	Val	Ser	Asp	Phe	Thr	Asp
		545			550					555					560
Ser	Val	Arg	Asp	Pro	Lys	Thr	Ser	Glu	Ile	Leu	Asp	Ile	Ser	Pro	Cys
				565					570					575	
Ser	Phe	Gly	Gly	Val	Ser	Val	Ile	Thr	Pro	Gly	Thr	Asn	Ala	Ser	Ser
				580				585					590		
Glu	Val	Ala	Val	Leu	Tyr	Gln	Asp	Val	Asn	Cys	Thr	Asp	Val	Ser	Thr
		595					600					605			
Ala	Ile	His	Ala	Asp	Gln	Leu	Thr	Pro	Ala	Trp	Arg	Ile	Tyr	Ser	Thr
		610				615					620				
Gly	Asn	Asn	Val	Phe	Gln	Thr	Gln	Ala	Gly	Cys	Leu	Ile	Gly	Ala	Glu
		625			630					635					640
His	Val	Asp	Thr	Ser	Tyr	Glu	Cys	Asp	Ile	Pro	Ile	Gly	Ala	Gly	Ile
				645					650					655	
Cys	Ala	Ser	Tyr	His	Thr	Val	Ser	Leu	Leu	Arg	Ser	Thr	Ser	Gln	Lys
				660				665						670	
Ser	Ile	Val	Ala	Tyr	Thr	Met	Ser	Leu	Gly	Ala	Asp	Ser	Ser	Ile	Ala
		675					680					685			
Tyr	Ser	Asn	Asn	Thr	Ile	Ala	Ile	Pro	Thr	Asn	Phe	Ser	Ile	Ser	Ile
		690				695					700				
Thr	Thr	Glu	Val	Met	Pro	Val	Ser	Met	Ala	Lys	Thr	Ser	Val	Asp	Cys
		705			710					715					720
Asn	Met	Tyr	Ile	Cys	Gly	Asp	Ser	Thr	Glu	Cys	Ala	Asn	Leu	Leu	Leu
				725					730					735	
Gln	Tyr	Gly	Ser	Phe	Cys	Thr	Gln	Leu	Asn	Arg	Ala	Leu	Ser	Gly	Ile
				740				745					750		
Ala	Ala	Glu	Gln	Asp	Arg	Asn	Thr	Arg	Glu	Val	Phe	Ala	Gln	Val	Lys
		755					760					765			

-continued

Gln Met Tyr Lys Thr Pro Thr Leu Lys Tyr Phe Gly Gly Phe Asn Phe
 770 775 780
 Ser Gln Ile Leu Pro Asp Pro Leu Lys Pro Thr Lys Arg Ser Phe Ile
 785 790 795 800
 Glu Asp Leu Leu Phe Asn Lys Val Thr Leu Ala Asp Ala Gly Phe Met
 805 810 815
 Lys Gln Tyr Gly Glu Cys Leu Gly Asp Ile Asn Ala Arg Asp Leu Ile
 820 825 830
 Cys Ala Gln Lys Phe Asn Gly Leu Thr Val Leu Pro Pro Leu Leu Thr
 835 840 845
 Asp Asp Met Ile Ala Ala Tyr Thr Ala Ala Leu Val Ser Gly Thr Ala
 850 855 860
 Thr Ala Gly Trp Thr Phe Gly Ala Gly Ala Ala Leu Gln Ile Pro Phe
 865 870 875 880
 Ala Met Gln Met Ala Tyr Arg Phe Asn Gly Ile Gly Val Thr Gln Asn
 885 890 895
 Val Leu Tyr Glu Asn Gln Lys Gln Ile Ala Asn Gln Phe Asn Lys Ala
 900 905 910
 Ile Ser Gln Ile Gln Glu Ser Leu Thr Thr Thr Ser Thr Ala Leu Gly
 915 920 925
 Lys Leu Gln Asp Val Val Asn Gln Asn Ala Gln Ala Leu Asn Thr Leu
 930 935 940
 Val Lys Gln Leu Ser Ser Asn Phe Gly Ala Ile Ser Ser Val Leu Asn
 945 950 955 960
 Asp Ile Leu Ser Arg Leu Asp Lys Val Glu Ala Glu Val Gln Ile Asp
 965 970 975
 Arg Leu Ile Thr Gly Arg Leu Gln Ser Leu Gln Thr Tyr Val Thr Gln
 980 985 990
 Gln Leu Ile Arg Ala Ala Glu Ile Arg Ala Ser Ala Asn Leu Ala Ala
 995 1000 1005
 Thr Lys Met Ser Glu Cys Val Leu Gly Gln Ser Lys Arg Val Asp
 1010 1015 1020
 Phe Cys Gly Lys Gly Tyr His Leu Met Ser Phe Pro Gln Ala Ala
 1025 1030 1035
 Pro His Gly Val Val Phe Leu His Val Thr Tyr Val Pro Ser Gln
 1040 1045 1050
 Glu Arg Asn Phe Thr Thr Ala Pro Ala Ile Cys His Glu Gly Lys
 1055 1060 1065
 Ala Tyr Phe Pro Arg Glu Gly Val Phe Val Phe Asn Gly Thr Ser
 1070 1075 1080
 Trp Phe Ile Thr Gln Arg Asn Phe Phe Ser Pro Gln Ile Ile Thr
 1085 1090 1095
 Thr Asp Asn Thr Phe Val Ser Gly Asn Cys Asp Val Val Ile Gly
 1100 1105 1110
 Ile Ile Asn Asn Thr Val Tyr Asp Pro Leu Gln Pro Glu Leu Asp
 1115 1120 1125
 Ser Phe Lys Glu Glu Leu Asp Lys Tyr Phe Lys Asn His Thr Ser
 1130 1135 1140
 Pro Asp Val Asp Leu Gly Asp Ile Ser Gly Ile Asn Ala Ser Val
 1145 1150 1155
 Val Asn Ile Gln Lys Glu Ile Asp Arg Leu Asn Glu Val Ala Lys
 1160 1165 1170

-continued

Asn Leu Asn Glu Ser Leu Ile Asp Leu Gln Glu Leu Gly Lys Tyr
 1175 1180 1185
 Glu Gln Tyr Ile Lys Trp Pro Trp Tyr Val Trp Leu Gly Phe Ile
 1190 1195 1200
 Ala Gly Leu Ile Ala Ile Val Met Val Thr Ile Leu Leu Cys Cys
 1205 1210 1215
 Met Thr Ser Cys Cys Ser Cys Leu Lys Gly Ala Cys Ser Cys Gly
 1220 1225 1230
 Ser Cys Cys Lys Phe Asp Glu Asp Asp Ser Glu Pro Val Leu Lys
 1235 1240 1245
 Gly Val Lys Leu His Tyr Thr
 1250 1255

<210> SEQ ID NO 5
 <211> LENGTH: 274
 <212> TYPE: PRT
 <213> ORGANISM: Coronavirus

<400> SEQUENCE: 5

Met Asp Leu Phe Met Arg Phe Phe Thr Leu Gly Ser Ile Thr Ala Gln
 1 5 10 15
 Pro Val Lys Ile Asp Asn Ala Ser Pro Ala Ser Thr Val His Ala Thr
 20 25 30
 Ala Thr Ile Pro Leu Gln Ala Ser Leu Pro Phe Gly Trp Leu Val Ile
 35 40 45
 Gly Val Ala Phe Leu Ala Val Phe Gln Ser Ala Thr Lys Ile Ile Ala
 50 55 60
 Leu Asn Lys Arg Trp Gln Leu Ala Leu Tyr Lys Gly Phe Gln Phe Ile
 65 70 75 80
 Cys Asn Leu Leu Leu Leu Phe Val Thr Ile Tyr Ser His Leu Leu Leu
 85 90 95
 Val Ala Ala Gly Met Glu Ala Gln Phe Leu Tyr Leu Tyr Ala Leu Ile
 100 105 110
 Tyr Phe Leu Gln Cys Ile Asn Ala Cys Arg Ile Ile Met Arg Cys Trp
 115 120 125
 Leu Cys Trp Lys Cys Lys Ser Lys Asn Pro Leu Leu Tyr Asp Ala Asn
 130 135 140
 Tyr Phe Val Cys Trp His Thr His Asn Tyr Asp Tyr Cys Ile Pro Tyr
 145 150 155 160
 Asn Ser Val Thr Asp Thr Ile Val Val Thr Glu Gly Asp Gly Ile Ser
 165 170 175
 Thr Pro Lys Leu Lys Glu Asp Tyr Gln Ile Gly Gly Tyr Ser Glu Asp
 180 185 190
 Arg His Ser Gly Val Lys Asp Tyr Val Val Val His Gly Tyr Phe Thr
 195 200 205
 Glu Val Tyr Tyr Gln Leu Glu Ser Thr Gln Ile Thr Thr Asp Thr Gly
 210 215 220
 Ile Glu Asn Ala Thr Phe Phe Ile Phe Asn Lys Leu Val Lys Asp Pro
 225 230 235 240
 Pro Asn Val Gln Ile His Thr Ile Asp Gly Ser Ser Gly Val Ala Asn
 245 250 255
 Pro Ala Met Asp Pro Ile Tyr Asp Glu Pro Thr Thr Thr Ser Val
 260 265 270
 Pro Leu

-continued

<210> SEQ ID NO 6
 <211> LENGTH: 154
 <212> TYPE: PRT
 <213> ORGANISM: Coronavirus

<400> SEQUENCE: 6

```

Met Met Pro Thr Thr Leu Phe Ala Gly Thr His Ile Thr Met Thr Thr
1          5          10          15
Val Tyr His Ile Thr Val Ser Gln Ile Gln Leu Ser Leu Leu Lys Val
          20          25          30
Thr Ala Phe Gln His Gln Asn Ser Lys Lys Thr Thr Lys Leu Val Val
          35          40          45
Ile Leu Arg Ile Gly Thr Gln Val Leu Lys Thr Met Ser Leu Tyr Met
          50          55          60
Ala Ile Ser Pro Lys Phe Thr Thr Ser Leu Ser Leu His Lys Leu Leu
          65          70          75          80
Gln Thr Leu Val Leu Lys Met Leu His Ser Ser Ser Leu Thr Ser Leu
          85          90          95
Leu Lys Thr His Arg Met Cys Lys Tyr Thr Gln Ser Thr Ala Leu Gln
          100          105          110
Glu Leu Leu Ile Gln Gln Trp Ile Gln Phe Met Met Ser Arg Arg Arg
          115          120          125
Leu Leu Ala Cys Leu Cys Lys His Lys Lys Val Ser Thr Asn Leu Cys
          130          135          140
Thr His Ser Phe Arg Lys Lys Gln Val Arg
          145          150

```

<210> SEQ ID NO 7
 <211> LENGTH: 76
 <212> TYPE: PRT
 <213> ORGANISM: Coronavirus

<400> SEQUENCE: 7

```

Met Tyr Ser Phe Val Ser Glu Glu Thr Gly Thr Leu Ile Val Asn Ser
1          5          10          15
Val Leu Leu Phe Leu Ala Phe Val Val Phe Leu Leu Val Thr Leu Ala
          20          25          30
Ile Leu Thr Ala Leu Arg Leu Cys Ala Tyr Cys Cys Asn Ile Val Asn
          35          40          45
Val Ser Leu Val Lys Pro Thr Val Tyr Val Tyr Ser Arg Val Lys Asn
          50          55          60
Leu Asn Ser Ser Glu Gly Val Pro Asp Leu Leu Val
          65          70          75

```

<210> SEQ ID NO 8
 <211> LENGTH: 221
 <212> TYPE: PRT
 <213> ORGANISM: Coronavirus

<400> SEQUENCE: 8

```

Met Ala Asp Asn Gly Thr Ile Thr Val Glu Glu Leu Lys Gln Leu Leu
1          5          10          15
Glu Gln Trp Asn Leu Val Ile Gly Phe Leu Phe Leu Ala Trp Ile Met
          20          25          30
Leu Leu Gln Phe Ala Tyr Ser Asn Arg Asn Arg Phe Leu Tyr Ile Ile
          35          40          45
Lys Leu Val Phe Leu Trp Leu Leu Trp Pro Val Thr Leu Ala Cys Phe

```

-continued

```

50           55           60
Val Leu Ala Ala Val Tyr Arg Ile Asn Trp Val Thr Gly Gly Ile Ala
65           70           75           80
Ile Ala Met Ala Cys Ile Val Gly Leu Met Trp Leu Ser Tyr Phe Val
85           90
Ala Ser Phe Arg Leu Phe Ala Arg Thr Arg Ser Met Trp Ser Phe Asn
100          105          110
Pro Glu Thr Asn Ile Leu Leu Asn Val Pro Leu Arg Gly Thr Ile Val
115          120          125
Thr Arg Pro Leu Met Glu Ser Glu Leu Val Ile Gly Ala Val Ile Ile
130          135          140
Arg Gly His Leu Arg Met Ala Gly His Pro Leu Gly Arg Cys Asp Ile
145          150          155          160
Lys Asp Leu Pro Lys Glu Ile Thr Val Ala Thr Ser Arg Thr Leu Ser
165          170          175
Tyr Tyr Lys Leu Gly Ala Ser Gln Arg Val Gly Thr Asp Ser Gly Phe
180          185          190
Ala Ala Tyr Asn Arg Tyr Arg Ile Gly Asn Tyr Lys Leu Asn Thr Asp
195          200          205
His Ala Gly Ser Asn Asp Asn Ile Ala Leu Leu Val Gln
210          215          220

```

<210> SEQ ID NO 9
 <211> LENGTH: 63
 <212> TYPE: PRT
 <213> ORGANISM: Coronavirus

<400> SEQUENCE: 9

```

Met Phe His Leu Val Asp Phe Gln Val Thr Ile Ala Glu Ile Leu Ile
1           5           10          15
Ile Ile Met Arg Thr Phe Arg Ile Ala Ile Trp Asn Leu Asp Val Ile
20          25          30
Ile Ser Ser Ile Val Arg Gln Leu Phe Lys Pro Leu Thr Lys Lys Asn
35          40          45
Tyr Ser Glu Leu Asp Asp Glu Pro Met Glu Leu Asp Tyr Pro
50          55          60

```

<210> SEQ ID NO 10
 <211> LENGTH: 122
 <212> TYPE: PRT
 <213> ORGANISM: Coronavirus

<400> SEQUENCE: 10

```

Met Lys Ile Ile Leu Phe Leu Thr Leu Ile Val Phe Thr Ser Cys Glu
1           5           10          15
Leu Tyr His Tyr Gln Glu Cys Val Arg Gly Thr Thr Val Leu Leu Lys
20          25          30
Glu Pro Cys Pro Ser Gly Thr Tyr Glu Gly Asn Ser Pro Phe His Pro
35          40          45
Leu Ala Asp Asn Lys Phe Ala Leu Thr Cys Thr Ser Thr His Phe Ala
50          55          60
Phe Ala Cys Ala Asp Gly Thr Arg His Thr Tyr Gln Leu Arg Ala Arg
65          70          75          80
Ser Val Ser Pro Lys Leu Phe Ile Arg Gln Glu Glu Val Gln Gln Glu
85          90          95
Leu Tyr Ser Pro Leu Phe Leu Ile Val Ala Ala Leu Val Phe Leu Ile

```


-continued

Leu Asp Arg Leu Asn Gln Leu Glu Ser Lys Val Ser Gly Lys Gly Gln
 225 230 235 240
 Gln Gln Gln Gly Gln Thr Val Thr Lys Lys Ser Ala Ala Glu Ala Ser
 245 250 255
 Lys Lys Pro Arg Gln Lys Arg Thr Ala Thr Lys Gln Tyr Asn Val Thr
 260 265 270
 Gln Ala Phe Gly Arg Arg Gly Pro Glu Gln Thr Gln Gly Asn Phe Gly
 275 280 285
 Asp Gln Asp Leu Ile Arg Gln Gly Thr Asp Tyr Lys His Trp Pro Gln
 290 295 300
 Ile Ala Gln Phe Ala Pro Ser Ala Ser Ala Phe Phe Gly Met Ser Arg
 305 310 315 320
 Ile Gly Met Glu Val Thr Pro Ser Gly Thr Trp Leu Thr Tyr His Gly
 325 330 335
 Ala Ile Lys Leu Asp Asp Lys Asp Pro Gln Phe Lys Asp Asn Val Ile
 340 345 350
 Leu Leu Asn Lys His Ile Asp Ala Tyr Lys Thr Phe Pro Pro Thr Glu
 355 360 365
 Pro Lys Lys Asp Lys Lys Lys Lys Thr Asp Glu Ala Gln Pro Leu Pro
 370 375 380
 Gln Arg Gln Lys Lys Gln Pro Thr Val Thr Leu Leu Pro Ala Ala Asp
 385 390 395 400
 Met Asp Asp Phe Ser Arg Gln Leu Gln Asn Ser Met Ser Gly Ala Ser
 405 410 415
 Ala Asp Ser Thr Gln Ala
 420

<210> SEQ ID NO 13
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic oligonucleotide.

<400> SEQUENCE: 13

ctaacatgct taggataatg g

21

<210> SEQ ID NO 14
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic oligonucleotide.

<400> SEQUENCE: 14

gcctctcttg ttcttgetcg c

21

<210> SEQ ID NO 15
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic oligonucleotide.

<400> SEQUENCE: 15

caggaagcg taaaactcat c

21

<210> SEQ ID NO 16
 <211> LENGTH: 22

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide.

<400> SEQUENCE: 16

catgtgtggc ggctcactat at 22

<210> SEQ ID NO 17
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide.

<400> SEQUENCE: 17

gacactatta gcataagcag ttgtagca 28

<210> SEQ ID NO 18
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide.

<400> SEQUENCE: 18

ttaaaccagg tggaacatca tccggtg 27

<210> SEQ ID NO 19
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide.

<400> SEQUENCE: 19

ggagccttga atacacccaa ag 22

<210> SEQ ID NO 20
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide.

<400> SEQUENCE: 20

gcacggtggc agcattg 17

<210> SEQ ID NO 21
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide.

<400> SEQUENCE: 21

ccacattggc acccgcaatc c 21

<210> SEQ ID NO 22
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide.

<400> SEQUENCE: 22

-continued

caaacattgg ccgcaaatt	19
<210> SEQ ID NO 23 <211> LENGTH: 21 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide. <400> SEQUENCE: 23	
caatgcgtga cattccaaag a	21
<210> SEQ ID NO 24 <211> LENGTH: 26 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide. <400> SEQUENCE: 24	
cacaatttgc tccaagtgcc tctgca	26
<210> SEQ ID NO 25 <211> LENGTH: 21 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide. <400> SEQUENCE: 25	
gaagtaccat ctggggctga g	21
<210> SEQ ID NO 26 <211> LENGTH: 18 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide. <400> SEQUENCE: 26	
ccgaagagct acccgacg	18
<210> SEQ ID NO 27 <211> LENGTH: 26 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide. <400> SEQUENCE: 27	
ctctttcatt ttgccgtcac caccac	26
<210> SEQ ID NO 28 <211> LENGTH: 24 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide. <400> SEQUENCE: 28	
agctctccct agcattattc actg	24
<210> SEQ ID NO 29 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence	

-continued

```

<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide.

<400> SEQUENCE: 29

caccacattt tcatcgaggc                                20

<210> SEQ ID NO 30
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide.

<400> SEQUENCE: 30

taccctcgat cgtactccgc gt                              22

<210> SEQ ID NO 31
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide.

<400> SEQUENCE: 31

tgtaggcact gattcaggtt ttg                             23

<210> SEQ ID NO 32
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide.

<400> SEQUENCE: 32

cggcgtggtc tgtatttaat tta                             23

<210> SEQ ID NO 33
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide.

<400> SEQUENCE: 33

ctgcatacaa ccgctaccgt attggaa                         27

<210> SEQ ID NO 34
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide.

<400> SEQUENCE: 34

gggttgggac taccctaagt gtga                            24

<210> SEQ ID NO 35
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (12)..(12)
<223> OTHER INFORMATION: "n" equals inosine.

```

-continued

<400> SEQUENCE: 35

taacacacaa cncctatc a

21

<210> SEQ ID NO 36

<211> LENGTH: 19

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide.

<400> SEQUENCE: 36

agatttgac ctgcgagcg

19

<210> SEQ ID NO 37

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide.

<400> SEQUENCE: 37

gagcggctgt ctccacaagt

20

<210> SEQ ID NO 38

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide.

<400> SEQUENCE: 38

ttctgacctg aaggctctgc gcg

23

The invention claimed is:

1. A method of detecting a severe acute respiratory syndrome-associated coronavirus (SARS-CoV) in a sample comprising:

contacting the sample with a pair of nucleic acid primers that hybridize to a SARS-CoV nucleic acid, wherein at least one primer is 5'-end labeled with a reporter dye, and wherein at least one of the primers comprises the sequence as set forth in any one of SEQ ID NOs: 13-15; amplifying the SARS-CoV nucleic acid or a fragment thereof from the sample utilizing the pair of nucleic acid primers;

electrophoresing the amplified products; and detecting the 5'-end labeled reporter dye, thereby detecting a SARS-CoV.

2. The method of claim 1, wherein the amplification utilizes reverse transcriptase-polymerase chain reaction.

3. A method of detecting a severe acute respiratory syndrome-associated coronavirus (SARS-CoV) in a sample, comprising:

contacting the sample with a pair of nucleic acid primers that hybridize to a SARS-CoV nucleic acid, wherein at least one of the nucleic acid primers comprises the sequence as set forth in any one of SEQ ID NOs: 13-15; amplifying the SARS-CoV nucleic acid or a fragment thereof from the sample utilizing the pair of nucleic acid primers;

adding to the amplified SARS-CoV nucleic acid or the fragment thereof a SARS-CoV probe that hybridizes to the SARS-CoV nucleic acid, wherein the SARS-CoV probe is labeled with a 5'-reporter dye and a 3'-quencher dye;

performing one or more additional rounds of amplification with Taq DNA polymerase; and

detecting fluorescence of the 5'-reporter dye, thereby detecting a SARS-CoV.

4. A kit for detecting a severe acute respiratory syndrome-associated coronavirus (SARS-CoV) in a sample, comprising a pair of nucleic acid primers that hybridize under stringent conditions to a SARS-CoV nucleic acid, wherein at least one of the primers comprises the sequence as set forth in any one of SEQ ID NOs: 13-15.

5. The kit of claim 4, wherein one primer is 5'-end labeled with a reporter dye.

6. The kit of claim 4, further comprising a SARS-CoV probe that hybridizes to the SARS-CoV nucleic acid amplified by the pair of primers, wherein the SARS-CoV probe is labeled with a 5'-reporter dye and a 3'-quencher dye.

7. The kit of claim 4, further comprising an isolated SARS-CoV organism.

* * * * *